



International Journal of Fauna and Biological Studies

Available online at www.faunajournal.com

I	International
J	Journal of
F	Fauna And
B	Biological
S	Studies

ISSN 2347-2677
IJFBS 2015; 2(4): 80-85
Received: 08-05-2015
Accepted: 11-06-2015

Muneesh kumar
Department of Zoology,
Govt. S.S.L. Jain P.G. College,
Vidisha, Barkatullah University
Bhopal (M.P.) India.

Parvinder kumar
Department of Zoology
University of Jammu, Jammu,
India.

Sangeeta Devi
Department of Botany,
Govt. S.S.L Jain PG College
Vidisha, Bhopal (M.P.)

Toxicity of Copper Sulphate on Behavioural Parameter and Respiratory Surveillance in Freshwater Catfish, *Clarias Batrachus* (Linn.)

Muneesh kumar, Parvinder kumar, Sangeeta Devi

Abstract

Industries are the major sources of heavy metal pollution and it is released into water and soil. Heavy metals cause several ill effects to aquatic living organisms and environment. The toxicity of copper sulphate to the freshwater fish, *Clarias batrachus* was studied using static bioassay method. The LC₅₀ in 96h was found to be 33 µg/L. Behavioural changes when exposed to lethal concentration of copper sulphate showed increased opercular movement, increased surface behaviour, loss of equilibrium, change in body colour, increased secretion of mucus, irregular swimming activity, rapid jerk movement, partial jerk and aggressiveness. The swimming behaviour was in a cork-screw palter, rotating along horizontal axis. In sub lethal treatment, the schooling behaviour of the fish was slowly disrupted, the ventilation rate was increased. The fish at 21st days of exposure exhibited balanced swimming and active feeding and behaved in normal way. Oxygen consumption was decreased in lethal concentration (-22.64 to -70.13%), but in sub lethal concentration decreased trend was improved and reached normal level at 21st day (-25.10 to -2.19). Alterations in oxygen consumption may be due to respiratory distress as a consequence of impairment in oxidative metabolism. Fish in sub lethal concentration were found under stress, but that was not fatal.

Keywords: Copper Sulphate, Toxicity, Behaviour, LC₅₀, *Clarias batrachus*, Oxygen consumption.

1. Introduction

The problem of pollution of the water where the wastes are usually discharged has increased to a great extent in recent years. Aquatic life is strongly influenced by physical properties of a water body. It is known that heavy metals as well as agro-pollutants are potentially harmful to the aquatic lives. All pesticides applied for the pest control eventually pollute the water resources either in their original chemical form or in some degraded variety. On the other hand, all industries discharge their effluents indiscriminately in the adjoining water areas and frequently cause serious hazards to aquatic life. Among the aqua fauna, fishes are affected to a significant extent. The contamination of inland & surface waters and land/soil, due to the release of variety of chemicals may prove toxic to all classes of living organisms. Among metals, copper is used in industries manufacturing organic chemicals, fertilizers, iron and steel works, electrical works, antifouling paints, pulp and paper industries, pesticides, fungicides and automobile accessories. Copper sulphate is widely used as an algaecide for controlling phytoplankton in fish ponds and lakes as well as a herbicide used in aquatic weed control since 1882 (Carbonell and Tarazona). Copper intoxication increases mortality of fish offspring and weakens their condition, resistance and viability (Jezierska and Witesca). Intensive industrial developments in the last few decades have increased the concentration of copper in aquatic ecosystems and that affected fish and deteriorate the natural resources. *Clarias batrachus*, one of the Indian major carp is an edible freshwater fish of great economic importance. In the present paper, the authors describe the impact of copper sulphate on behavioural changes and oxygen consumption of the said species.

2. Materials and Methods

Adult and live fish *Clarias batrachus* were collected from the farm Patra and Bhadbhada Bhopal M.P.) brought to the laboratory, cleaned by using 0.1% KMnO₄ to avoid dermal infection. Only healthy fishes (Length: 12-15cm, Weight: 50-60g) were taken for experiment. Fishes were acclimatized in glass aquaria for 15 days and were fed with fish food

Correspondence:

Muneesh kumar
Department of Zoology Govt.
S.S.L. Jain P.G. College, Vidisha,
Barkatullah University Bhopal
(M.P.) India.

(earthworms) and water in the aquaria was replaced by freshwater at every 24h.

2.1. Physico-chemical parameters of water

The physico-chemical parameters of the water used for fish bioassay were carried out according to the methods described in Standard Methods (APHA-AWWA-WEF, 1998) [2]. The water quality parameters were as follows: temperature $28\pm1^{\circ}\text{C}$, salinity 198 mg/L, pH 8, dissolved oxygen 6.7 to 7.2 mg/L, chloride 46.3 mg/L, sodium 1.22 mg/L, potassium 30.5 mg/L, calcium 17.04 mg/L, magnesium 1 mg/L, carbon dioxide 9.0 mg/L, Hardness 115 mg/L, oxygen per cent<R> saturation 57 mg/L (as CaCO_3) and specific gravity 1.00374.

2.2. Toxicant selected and preparation of stock solution

The present study, copper sulphate (CuSO_4) was used as a toxicant. The stock solution of copper (1000mg/L) was prepared according to the method prescribed in the Standard Methods (APHA-AWWA-WEF, 1998) [2]. For experimental purpose, the required copper concentration was drawn from the prepared stock solution.

2.3. Toxicity evaluation

The percent mortality of fish in different concentrations of copper sulphate was determined at 96h exposure. For this, the experimental fish were divided into batches of ten each, and were exposed to different concentrations of copper sulphate ranging from 10 $\mu\text{g}/\text{L}$ to 55 $\mu\text{g}/\text{L}$ (i.e. 0.010 to 0.05 mg/L). This range was obtained on trial and error basis. Toxicity evaluation was carried out in static water (Doudoroff *et al.*, 1951) [10] and mortality rate was observed and recorded for all the concentration after 96 hours. A batch of fish was also maintained simultaneously in freshwater medium without copper, which served as negative control. All the experiments were performed in duplicates and repeated thrice to confirm the results. The mean values were derived following the method of Finney Probit Kill Theory (1971) [13] and Dragstedt

and Behren's equation (Carpenter, 1982) [5]. For long-term effect one tenth of LC_{50} copper sulphate was taken in the present study copper sulphate.

2.4. Fixation of exposure periods

In order to understand the influence of time over toxicity effect of lethal concentration of copper sulphate on *Clarias batrachus* was studied at different periods of exposure. Before experimentation, healthy fishes were collected from the large cement tank with the help of nylon net and hand net. They were acclimatized to laboratory conditions in glass troughs for fifteen days. Each trough contained 15 L of water with uniform sized fish (length 12-15 cm; weight 50-60 g). They were fed with commercial fish food pellets during acclimation. After 15 days, if fishes were in normal behavioural activity and good health conditions, those species were selected for experiment purpose. The fish were divided into two groups. One group without copper sulphate served as control and the other group was exposed to lethal concentration of copper sulphate for 1, 2, 3 and 4 days were chosen to observe the short-term effects and for sub lethal concentration of copper sulphate for 1, 7, 14 and 21 days were chosen to observe the long-term effects. During this experiment the behavioural changes were critically observed.

3. Results

3.1. Toxicity Studies: The percentage mortality of *Clarias batrachus* was observed to be 0% and 100% at copper sulphate concentration of 10 $\mu\text{g}/\text{L}$ and 55 $\mu\text{g}/\text{L}$ (Table 1). The LC_{50} value obtained through sigmoid curve was 33 $\mu\text{g}/\text{L}$ and linear curve was found to be 32 $\mu\text{g}/\text{L}$. The LC_{50} value obtained were verified using Dragstedt and Behren's equation and was found to be 34 $\mu\text{g}/\text{L}$. Thus the average LC_{50} for 96 h was found to be 33 $\mu\text{g}/\text{L}$.

Table 1: Toxicity of copper sulphate on *Clarias batrachus*.

Sl. No.	Conc. of toxicant	Log conc. of toxicant	Number fish exposed	Alive	Dead	% Mortality	Probit Mortality
1.	10	1.0000	10	10	0	0	0.00
2.	15	1.1761	10	9	1	10	3.27
3.	20	1.3010	10	8	2	20	4.16
4.	25	1.3979	10	7	3	30	4.48
5.	30	1.4771	10	6	4	40	4.75
6.	35	1.5441	10	4	6	60	5.25
7.	40	1.6021	10	3	7	70	5.52
8.	45	1.6532	10	2	8	80	5.84
9.	50	1.6990	10	1	9	90	6.28
10.	55	1.7719	10	0	10	100	8.09

3.2. Behavioural observations Control fish

Normal fish maintained a fairly compact school, covering about one third of the bottom during the first seven days of the 21 days experiment. By seventh day, the school became less compact covering up to two third of the tank area. Fishes were observed to scrap the bottom surface. When startled, they instantly formed a tight school that was maintained briefly. They were sensitive to light and moved to the bottom of the tank when light was passed into the tank. Except a less response to form a dense school towards the end of the study, no other extraordinary behaviour was observed.

3.3. Exposed fish

The behavioural changes were observed when exposed to copper sulphate. The fish showed erratic swimming, hyper and

hypoxic, imbalance in posture, increase in surfacing activity, opercular movement, gradual loss in equilibrium and spreading of excess of mucus all over the surface of the body. The fish occupied twice the area than that of the control group. They were spread out and appeared to be swimming independent of one another. The swimming behaviour was in a cork-screw pattern, rotating along horizontal axis and followed by 'S' jerk, partial jerk, sudden rapid, non-directed spurt of forward movement. The fish eventually died with their mouth and opercular wide opened. A change in colour of gill lamellae from reddish to light brown with coagulation of mucus on the gill lamellae was seen in dead fish. In sub lethal treatment, the schooling behaviour of the fish was slowly disrupted during the first day. The ventilation rate was increased, hyperactivity, excitement, hyperventilation etc, were not much influenced on

exposure to the sub lethal concentration of copper sulphate at 7 and 14 days. Further, the fish at 21 days of exposure exhibited balanced swimming and active feeding. The fish behaved in normal way.

3.4. Oxygen consumption

The rate of whole animal oxygen consumption of control and copper sulphate treated fishes are presented in (Table 2). The data indicates that in fish exposed to lethal and sub lethal concentrations of copper sulphate, oxygen consumption was reduced. The whole animal O₂ consumption is reduced by

about 22.64 per cent on day one and reached maximum reduction of 70.13 per cent by day 4. Even during day 2 and 3 there was high rate of decrement as seen in (Table 2). The decrement was a sudden reduction from day one to day two, from then it was gradual reduction. However in sub lethal concentrations, the day 1 showed high rate of decrease of 25.10 per cent, which reached 19.22 by day 7. Subsequently, there was an improvement in O₂ consumption as by day 14 it was reduced to 12.92 per cent and day 21 showed minimal decrease which was recorded at 2.19 per cent. All the results are significant at 0.05, levels.

Table 2: Oxygen consumption of *Clarias batrachus* exposed to copper sulphate

	Control	1	2	3	4	1	7	14	21
Mean	0.8234	0.6534	0.5557	0.4433	0.2459	0.6167	0.6651	0.7170	0.8053
% change		-20.6440	-32.5123	-46.1677	-70.1309	-25.1067	-19.2232	-12.9288	-2.1979
SD	0.0030	0.0011	0.0018	0.0011	0.0013	0.0018	0.0020	0.0010	0.0042

4. Discussion

In the science of aquatic toxicology, fish play an important role in toxicity testing and hazard evaluation, as do the white rat and guinea pig in mammalian toxicology (Annon, 1972) [1]. The assessment of toxicity of copper sulphate with reference to aquatic biota, especially fish is crucial in establishing the toxicity evaluation. The LC₅₀ 96 hrs value of copper sulphate to *Clarias batrachus* was found to be 33 µg/l (Table 1). This data clearly indicates that copper sulphate is also toxic to *Clarias batrachus* as to other fish species mentioned in the table, copper sulphate is known to be readily taken up by aquatic organisms and bioconcentration factors ranged from 130 to 4,900 for various organisms (Snail, Daphnia and Fish). In laboratory tests copper sulphate was reported to be highly toxic for aquatic organisms with LC₅₀ values ranging from 0.02 µg/l to 1 µg/l for newly hatched shrimp. Various symptoms of poisoning can be observed from studies involving the determination of LC₅₀. In the present study, the fish maintained in normal freshwater behaved in usual manner i.e., they were very active with their well-coordinated movements. They were alert at slightest disturbance. But at the sub lethal concentrations of copper sulphate they became irritable and hyper-excited. Jumping movements as well as restlessness were observed and finally the fish turned upside down. Mucus secretion and loss of equilibrium were also observed. They slowly became sluggish with short jerky movements, surfacing and gulping of air and erratic circular movements. Finally they settled down at the bottom with loss of equilibrium and rolling of the body, convulsions prior to death. The fish very often come to the surface in order to avoid toxic environment. Moreover, examination of the gill of dead fish revealed that the gill lamellae colour was changed from red to brown. *Tilapia* exposed to lethal and sub lethal concentration of endosulfan and lindane exhibited abnormal behaviour at lethal concentration; a sudden heavy stress on the fish showed erratic swimming, convulsion, spiralling, tremors, jerky movements and rapid opercular movements. The fish struggled hard for breathing, often moved to the surface to engulf atmospheric air and tried to escape the toxic aquatic medium. After a few hours, equilibrium was lost and the fishes spiralled and slowly moved upward in a vertical position. Finally they lost equilibrium completely and were flat at the bottom (Thorat, 2001) [38]. Similar symptoms were also observed by Deva Prakasa Raju, 2000 [9] in *L. rohita* and Prashanth and Neelgund 2008 [26] in *C. mrigala* respectively. Some workers have reported the toxicity evaluation of

cyanides in the freshwater fish, *Catla catla* (Prashanth and Patil 2006) [25]. The difference might be due to either potency of toxicant or difference in the test conditions. The mean LC₅₀ value of copper sulphate estimated in the present investigation was 33µg/l (Table 1). The unusual behaviour of the fish, *Clarias batrachus* in stress condition may be due to obstructed functions of neurotransmitters. The gill opercular movements increased initially to support enhanced physiological activities in stressful habitat and later decreased may be due to mucus accumulation of gill. The toxic stress of pesticides has direct bearing on tissue chemical compounds (Tilak and Yacobu, 2002) [39]. This was also reported by David, 1995 [7] and Muniyan and Veeraraghavan, 1999 [19], Chaudhary *et al.*, 2001 [6]. The excessive secretion of mucus over the gills may inhibit the diffusion of oxygen during the process of gaseous exchange. It suggests that the copper sulphate is not safe to non-target organisms like fishes. In order to protect whole aquatic ecosystem, awareness must be given to the farmers to control the agricultural pests by biological methods.

The behavioural changes are the manifestation of motivational, biochemical, physiological and environmentally influenced state of the organism. The migration of fish to the bottom of the tank following the addition of copper sulphate clearly indicates the avoidance behaviour of the fish, which was reported by Murthy (1987) [20] in trout. The opercular movement of the fish ceases immediately following exposure to copper. The increase in opercular movement and corresponding increase in frequency of surfacing of fish clearly indicates that fish adaptively shifts towards aerial respiration (by obtaining atmospheric oxygen surfacing) and the fish tries to avoid contact with the copper through gill chamber (Santhakumar and Balaji, 2000; Prashanth & Patil, 2006) [31, 32, 25]. The increased ventilation rate by rapid, repeated opening and closing of mouth and opercular coverings accompanied by partially extended fins (caughing) was observed in the present study. This could be due to clearance of the accumulated mucus debris in the gill region for proper breathing, which was suggested by Prashanth *et al.*, 2005 [27]. The erratic swimming of the treated fish indicates loss of equilibrium.

It is likely that the region in the brain, which is associated with the maintenance of equilibrium, should have been affected (Deva Parkasa Raju, 2000; Prashanth *et al.*, 2005; Prashanth & Patil, 2006) [9, 27, 25]. The erratic swimming, jerky movements and convulsions before death were evident and the serenity varied with pesticide concentration. It indicates the sings of

asphyxiation as indicated by gasping to death when fish, *Sarotherodon mossambicus* exposed dimethorate (Kalavathy et al., 2001) [16]. The surfacing phenomenon of fish observed under copper exposure might either be due to hypoxic condition of the fish as reported by Radhaiah and Jayantha Rao (1988) [28]. This fact was clearly evidenced in the present study. The observation on the metabolic shift from aerobic to an anaerobic condition involving glycolytic oxidation with enormous amount of lactic acid accumulation were also seen. Chronic exposure of finfish to aroclor was found to induce surfacing phenomenon of fish as pointed out by Drummond et al., 1986 [11]. Aggressive behaviour such as nudge and nip were increased following exposure to the toxic material. Orientation and locomotor patterns were found to be involved in most aspects of fish behaviour such as migration, mating, courtship and feeding, which were altered under stress conditions of environmental toxicants (David, 1995; Madhab Prasad et al., 2002; Prashanth et al., 2005; Prashanth & Patil, 2006) [7, 17, 27, 25]. The hyperexcitability of the fish invariably in the lethal exposure to free copper/pesticides may probably be due to the hindrance in the functioning of the enzyme AChE in relation to nervous system as suggested by many authors (Deva Prakasa Raju, 2000; Prashanth, 2003) [9, 23]. It leads to accumulation of acetylcholine, which is likely to cause prolonged excitatory post synaptic potential. This may first lead to stimulation and later cause a block in the cholinergic system.

The accumulation and increased secretion of mucus in the fish exposed to copper may be adaptive responses perhaps providing additional protection against corrosive nature of the pesticide and to avoid the absorptions of the toxicant by the general body surface. This agrees to the earlier findings done by David (1995) [7]. In the present study as evidenced by the results the abnormal changes in the fish exposed to lethal concentration free copper is time dependent. Since most fish breathe water in which they live, changes in the chemical properties thereof may be reflected in the animal's ventilator activity, particularly if the environment affects respiratory gas exchange (Sellers et al., 1975) [34]. Toxicants from the environment mainly enter fish by means of their respiratory systems (Tovell et al., 1975) [40]. A mechanism of toxicant uptake through gills probably occurs through pores by simple diffusion and is then absorbed through cell membranes (Opperhuizen et al., 1985) [21]. Studies on the course of oxygen consumption in lethal and sub lethal concentration indicate the sequence of the type of compensatory mechanism, if any, which operates within the animal to overcome the load of toxic stress.

From the results, presented in the table 2, it is clearly evident that copper sulphate affects the oxygen consumption of the fish, *Clarias batrachus* under lethal and sub lethal concentrations. The observed decrease in oxygen consumption by the whole animal may be due to the respiratory distress as a consequence of the impairment of oxidative metabolism. Several authors reported similar decline in whole animal oxygen consumption in different species of fishes exposed to toxicants (Kabeer et al., 1981; Rangaswamy, 1984; Deva Prakasa Raju, 2000; David et al., 2002) [15, 29, 9, 8]. Gills are the major respiratory organs and all metabolic pathways depend upon the efficiency of the gill for their energy supply and damage to these vital organs causes a chain of destructive events, which ultimately lead to respiratory distress (Radhaiah and Jayantha Rao, 1988; Esther, et al., 2001) [28, 12]. In consonance with this, he also reported that the depletion in O₂

consumption was due to the disorganization of the respiratory function caused by rupture in the respiratory epithelium of the gill. It is also due to the disturbance in mitochondrial integrity and decreased activities of some mitochondrial enzymes (Ravinder, 1988) [30]. Magare and Patil (2000) [18] reported a decrease in the rate of O₂ consumption in *Puntius ticto* exposed to endosulfan. In addition to gill damage decrease in haemoglobin content and decrease in tissues respiration (Sarkar, 1999) [33] may also interfere with respiratory process resulting in respiratory failure.

The decrement can also be attributed to the induction of hypoxic conditions within the animal due to the in-time contact of the respiratory surface with the polluted water resulting in the alteration of normal respiratory area of the animal. The secretion of mucus layer over the gill lamellae has been observed during copper stress. Excessive secretion of mucus over the gills may inhibit the diffusion of oxygen during the process of gaseous exchange (Muniyan and Veeragahavan, 1999; David et al., 2002) [19, 8]. The coagulation of mucus on the gills caused demolition of various important processes such as gas exchange, nitrogen excretion, salt balance and circulation of blood (Skidmore, 1964) [36]. The alternative reason for the decrease in the oxygen consumption would be due to the internal action of copper sulphate. This toxic substance appears to alter the metabolic cycle at subcellular level. Greater decrease in the rate of O₂ consumption of the fish, exposed to lethal concentration than the sub lethal concentration, may be due to the considerable damage to the gill structure and also due to the greater precipitation of mucus upon gill filaments leading to the clogging of gills.

Probably suffocation imposed by the coagulated mucus film and necrosis on the epithelial and inter lamellar cells of gills is one of the reasons for the death of animal in lethal concentration. Greater decrease in the rate of O₂ consumption of the fish in lethal concentration may also be due to the greater damage caused to R.B.C. as evident by the drastic decrease in the number of these cells (Venkataramana, 1987) [41]. Lowering of O₂ consumption of fish in the sub lethal concentration of copper sulphate may be mostly due to the lowering down of energy requirements and if so, such lowering of maintenance energy requirement is to be considered adaptive and even strategic. This lowering of maintenance energy requirements may be achieved by reducing osmotic gradient through the lowering of electrolyte levels in the body fluids. Further, there is evidence for a considerable metabolic reorganization and increased utilization of anaerobic metabolism in fishes exposed to sub lethal concentration of copper sulphate. If so, the lowering of the oxidative metabolism in *Clarias batrachus* might have been compensated at least by some degree of glycolysis. Behavioural characteristics are obviously sensitive indicators of toxicant effect. It is necessary, however, to select behavioural indices of monitoring that relate to the organisms behaviour in the field in order to derive a more accurate assessment of the hazards that a contaminant may pose in natural system, should be considered for species forming social organizational.

If social interactions are not considered, only a certain portion of a population may be protected, and the toxicity of contaminant may be underestimated. And also primarily influence modulations in the level of oxygen consumption. This has led to the imbalance in cellular homeostasis. The inhibition of oxygen consumption is explained by this manner

in which copper sulphate is incorporated into this fish system for energy. It is concluded that copper sulphate is innately to fish, *Clarias batrachus*, but many factors influence the degree of hazard. The above-analogy also warrants for an indispensable need to evaluate more toxicity data for wide range of animal groups of the eco-web in order to understand the broad spectrum of copper sulphate in comparison to other toxicants available. This also provides a platform to establish tolerable limits and safe levels of toxic agents for the biota of aquatic environment and to save the residue imbalance in aquatic bio-ecological cycles, which help in involving bio-detector monitoring.

5. Conclusion

After the above discussion it had been concluded that copper sulphate causes deleterious effects on fishes and much alters the behavioural and O₂ consumption characteristic. In sub lethal concentration it may not be fatal for an individual organism but it does affect the growth rate and reproduction resulting in disturbance to whole community and trophic levels of food chains, ultimately the ecosystem.

6. Acknowledgement

Authors are beholden to Principal Dr. A.K. Ganguly S.L.L. Jain P.G. College Vidisha, Bhopal M.P. India for providing laboratory facilities for this research work. This paper forms the part of Ph.D. thesis submitted by author to Barkatullah University, Bhopal M.P. (India).

7. References

1. Annon. EPA bans most DDT uses, readies lead action. Environ. Sci. Technol 1972; 6:675.
2. APHA/AWWA-WEF. Standard Methods for the examination of water and waste water. 20th Edn. American Public Health Association, Washington, DC, 1998.
3. Billard R, Roubaud P. The effect of metals and cyanide on fertilization in rainbow trout (*Salmo gairdneri*). Water Research 1985; 19:209-214.
4. Carbonell GJ, Tarazona V. A proposed method to diagnose acute copper poisoning in cultured rainbow trout, *Oncorhynchus mykiss*. Sci. of Total Environ 1993; 2:1329-1334.
5. Carpenter PL. Immunology and Serology, 3rd ed., W. B. Saunders Co., Philadelphia, USA, 1982.
6. Chaudhary, Varslia Rathod, Vankhede GN. Effect of water extract of the bark of *Buchanania langsdorffii* (Linn). On behaviour and Chromatophores of a freshwater fish, *Labeo rohita*. J Environ. Biol. 2001; 22(3):229-331.
7. David M. Effect of fenvalerate on behavioral, physiological and biochemical aspects of freshwater fish, *Labeo rohita* (Hamilton) Ph.D. Thesis S. K. University. Anantapur, A.P. India, 1995.
8. David M, Mushigeri SB, Prashanth MS. Toxicity of Fenvalerate to the Freshwater Fish, *Labeo rohita*. Geobios 2002; 29(1):25-27.
9. Deva Parkasa Raju B. Fenvalerate induced changes in protein metabolism of fresh water fish, *Tilapia mossambica*. Ph. D. Thesis, S. K. University. Anantapur, A.P. India, 2000.
10. Doudoroff P, Anderson BG, Burchick GE, Galtsolf BS, Hart NB, Patrick R *et al.* Bio-assay method for the evaluation of acute toxicity of industrial wastes to fish sewage. Ind. Wastes 1951; 23:1380-1397.
11. Drummond RA, Russom CL, Diannel Geiger, David L Defoe. Behavioural and morphological changes in freshwater minnow *Pimephales promelas*. As diagnostic end point for screening chemicals according to mode of action. Aqua. Toxicol 1986; 9:415-434.
12. Esther Joice P, Mallikaraj D, Parthi N, Natarajan GM, Sasikala G, Tamilselvi G *et al.* The effect of hypoxic stress on the bimodal respiration of *Macropodus cupanus*. J Eco. Res. Biocon. 2001; 2(1-2):76-80.
13. Finney DT. Probit Analysis. 3rd ed. Cambridge University Press. London, 1971.
14. Jeziorska B, Witesca M. Summary of metal induced disturbances in fish organism, Metal Toxicity to Fish. Wydawnictwo Akademii Podlaskiej, Siedlce, 2001, 214-243.
15. Kabeer Ahmed I, Jagannatha Rao KS, Ramana Rao KV. Effect of malathion exposure on some physiological parameters of whole blood and on tissue cations of teleost, *Tilapia mossambica*. J Biol. Sci. 1981; 3:17-21.
16. Kalavathy K, Sivakumari AA, Rashmi Chandran. Toxic effect of the pesticide dimethoate on the fish, *Sarotherodon mossambicus*. J Eco. Res. Bicon. 2001; 2(152):27-32.
17. Madhab Prades, Bandyopadhyay, Ajit Kumar Adithya. Xenobiotic impact on sensitivity in *Anabas testudineus* (both). J Ecobiol. 2002; 14(2):117-124.
18. Magare SR, Patil HT. Effect of pesticides on oxygen consumption. Red blood cell count and metabolites of fish, *Puntius ticto*. Enviro and Eco. 2000; 18(4):891-894.
19. Muniyan M, Veeraraghavan K. Acute toxicity of ethofenprox to the fresh water fish, *Oreochromis mossambicus* (Perers). J Environ. Biol. 1999; 20(1):153-155.
20. Murthy AS. Sub lethal effects of pesticides on fish. Toxicity of pesticide to fish. C. R. S. Press Boca Roton F. I. USA 1987; 4:55-100.
21. Opperhuizen A, Van Der Velde EW Gobas, FAPC Liem, DAK, Van Der Steen JMD. Relationship between bio-concentration in fish and steric factor of hydrophobic chemicals. Chemosphere 1985; 14(11-12):1871-1896.
22. Patil YB, Paknikar KM. Development of a process for detoxification of metal-cyanides from waste waters. Process Biochemistry 2000; 35:1139-1151.
23. Prashanth MS. Cypermethrin induced physiological, biochemical and histopathological, 2003.
24. Changes in freshwater fish, *Cirrhinus mrigala*. Ph.D., thesis, Karnataka University, Dharwad. India.
25. Prashanth MS, Yogesh B, Patil. Behavioural surveillance of Indian major carp *Catla Catla* (Hamilton) exposed to free cyanide. J of Cur. Sci. 2006; 9(1):313-318.
26. Prashanth MS, Neelgund SE. Free cyanideinduced Biochemical changes in Nitrogen metabolism of the Indian major carp, *Cirrhinus mrigala*. Journal of Basic and Clinical Physiology and Pharmacology. 2008; 18(4):277-287.
27. Prashanth MS, David M, Mathad SG. Behavioural Changes In Freshwater Fish, *Cirrhinus Mrigala* (Hamilton) Exposed To Cypermethrin. J of Envio. Bio. 2005; 26(13):141-144.
28. Radhaiah V, Jayantha Rao K. Behavioural response of fish, *Tilapia mossambica* exposed to fenvalerate. Environ. Ecol 1988; 6(2):2-23.
29. Rangaswamy CP. Impact of endosulfan toxicity on some physiological properties of the blood and aspects of

- energy metabolism of a fresh water fish, *Tilapia mossambica*. Ph.D thesis, S.V. University Tirurpathi, India, 1984.
30. Ravinder V. In vivo effects of decis on certain aspects of metabolism of freshwater fish, *Clarias batrachus*. Ph.D., Thesis, Osmania University, Hyderabad, India, 1988.
 31. Santhakumar M, Balaji M. Acute toxicity of an organophosphorus insecticide monocrotophos and its effects on behaviour of an air-breathing fish, *Anabas testudineus* (Bloch). J Environ. Biol. 2000; 21(2):121-123.
 32. Santhakumar M, Balaji M, Saravanan KR, Soumady D, Ramudu K. Effect of monocrotophos on the optomotor behaviour of an air breathing fish, *Anabas testudinarius* (Bloch). J Environ. Biol. 2000; 21(1):65-68.
 33. Sarkar SK. Effects of heavy metals (Copper sulphate and Cadmium sulphate) on the O₂ consumption of fish, *Cyprinus carpio*. U.P. J Zool. 1999; 19(1):13-16.
 34. Sellers CM (Jr), Health AG ML. The effects of sublethal concentration of copper and zinc on ventilatory activities, blood O₂ and pH in rainbow trout. *Salmo gairdneri*. Water. Res 1975; 99:401-408.
 35. Sharpe AG. The Chemistry of Cyano Complex and the Transition Metals. London: Academic Press, 1976.
 36. Skidmore JF. Toxicity of Zinc compounds to aquatic animals, with special reference to fish. Quarterly Rev. Bio 1964; 39(3):227-248.
 37. Solomonson LP, Spehar AM. Glyoxylate and cyanide formation. In: Cyanide in Biology (Edited by Vennesland, B., Conn, E.E., Knowles, C.J. and Wissing, F.), Academic Press, London, 1981, 363-370.
 38. Thorat SR. Chronic effect of endosulfan on freshwater fish, *Catla catla*, J Ecotoxicol. Envion. Moint. 2001; 11(4):221-223.
 39. Tilak KS, Yaeobu K. Toxicity and effect of fenvalerate on fish, *Ctenopharyngodon idellus*. J Ecotoxi, Envron. Moint. 2002; 12(1):09-15.
 40. Tovell PW A, Howes D, Newsome CS. Absorption metabolism and excretion by the gold fish of the anionic detergent, Sodium lauryl sulphate. Toxicology 1975; 6:17-29. Vanden Bercken J Vijverberg HPM, 1983.
 41. Venkataramana P. Studies on lethal and sub lethal toxicity of copper in the fresh water teleost, *Labeo rohita*. Ph.D. Thesis. S.K.University. Anantapur, Andhra Pradesh, India, 1987.