



ISSN 2347-2677

IJFBS 2017; 4(3): 133-138

Received: 27-03-2017

Accepted: 29-04-2017

**Anju Bharti**

Apart of Ph.D. Thesis of  
Research Scholar, Biopesticide  
and Toxicological Lab.,  
Department of Zoology, D.B.S.  
College, affiliated to CSJM  
University, Kanpur, India

**BS Chandel**

Biopesticide and Toxicological  
Lab., Department of Zoology,  
D.B.S. College, affiliated to  
CSJM University, Kanpur, India

## Insectifuge Biopotency of *Chrysanthemum cinerariifolium*, *Cichorium intybus*, *Inula racemosa* and *Tagetes minuta* extractives against okra aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae)

**Anju Bharti and BS Chandel**

### Abstract

In the present investigation alcoholic extract of ten indigenous naturally occurring asteraceous plants viz., aerial parts of *Acmella paniculata* (Wall ex DC.) R.K.Jansen, *Cichorium intybus* (L.), *Chromolaena odorata* Linn., *Chrysanthemum cinerariifolium* (trev.) Vis., *Inula racemosa* Hook. F., *Mantisalca duriaeri* Birq. Et Cavill., *Rechardia tingitana* (L.) Roth, *Rhaponticum acaule* (L.) DC, *Scorzonera undulate* Vahl, and *Tagetes minuta* Linn. Extractives were prepared under the laboratory conditions. For testing the repellent effect on okra leaves were used as food against nymph and adults of *Aphis gossypii* Glover. After four hours nymph and adults of *A. gossypii* were release and the data was collected on the number of nymph and adults of *A. gossypii* reached and repelled at each treated food. All the comparisons were made with control. It is evident from the results of repellent test against nymph and adults of *Aphis gossypii* Glover that all the asteraceous extractives showed a good repellent property, when compared with the control. The repellent properties were observed based on their minimum EC<sub>50</sub> values as: *C. intybus* (0.1359) > *I. racemosa* (0.1531) > *T. minuta* (0.1833) > *M. duriaeri* (0.3208) > *S. undulate* (0.4123) > *A. paniculata*, (0.6365) > *R. acaule* (0.6469) > *R. tingitana* (0.7124) > *C. odorata* (0.9025), respectively. The relative EC<sub>50</sub> values of the above extracts are in descending order, 6.6409 > 5.8948 > 4.9236 > 2.8132 > 1.4179 > 1.3951 > 1.2668 > 1.2161 > 1.1668 > 1.0000 times as repellents, whereas *A. paniculata* was taken as a unit. *C. intybus* and *I. racemosa* extract significantly reduced the population of treated nymphs and adults of *Aphis gossypii* on okra vegetable compared to the untreated control.

**Keywords:** *Aphis gossypii*, *Cichorium intybus*, *Inula racemosa*, *Tagetes minuta*, okra

### 1. Introduction

Bhindi, *Abelmoschus esculentus* (Linn.) Moench commonly known as okra is an important vegetable crop grown all over Indian sub Zoogeographical region (Radake and Undirwade 1981) [1]. Okra contains large quantities of vitamin B, vitamin C, carbohydrate, potassium, protein, folic acid, and calcium (Adeboye and Oputa, 1996) [2]. It's low in calories and has high dietary fiber content. Okra is a nutritious vegetable which plays an important role to meet the demand of vegetables of the India when vegetable are scanty in the market. Okra mucilage is suitable for medicinal and industrial application (Kumar *et al.* 2013) [3].

The okra aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae) is a serious pest of vegetables ie; okra, brinjal, pepper melon, cucumber, pumpkin and cotton. The crop is vulnerable to attack by many serious insect pests among them, okra aphid, *A. gossypii* is the most important pest causing direct damage to the marketable fruits (Konar and Rai, 1990) [4]. *Aphis gossypii* and *Earias vittella* is alone reported to cause 57.1% fruit infestation and 54.04% yield loss in okra, *Abelmoschus esculentus* (Chaudhary and Dadheech, 1989) [5]. Initially, leaves will yellow and with an increasing number of okra aphids, they will begin to curl. Leaves can be damaged to such an extent that they wilt and fall off. This impairs photosynthesis, thereby weakening the okra plant even further, and may render fruits unmarketable and human consumption.

No doubt the synthetic chemical insecticides are very effective and used in insect pest management for a long time and to ensure higher vegetable and crop production (Chandel *et al.* 2018) [6]. The injudicious and excessive use of these chemical pesticides led to many problems like development of resistance, environment pollutions and human health (Park *et al.* 2011) [7]. The side effects have forced to look for naturally occurring ecofriendly indigenous

### Correspondence

**Anju Bharti**

Apart of Ph.D. Thesis of  
Research Scholar, Biopesticide  
and Toxicological Lab.,  
Department of Zoology, D.B.S.  
College, affiliated to CSJM  
University, Kanpur, India

herbal alternatives aspect to chemical pesticides especially for vegetables like okra, where fruits are plucked at an interval of every 2-3 days (Chandel *et al.* 2018) <sup>[8]</sup>. Various plant extractives have been studied for insecticidal activity globally (Bajpai and Chandel, 2009, 2010 <sup>[9, 10]</sup> and majority of them are insect feeding deterrent (Antonius and Hagazy 1987, Chandel and Sengar 2018) <sup>[11, 12]</sup>. More than 140 compounds, which are chemically diverse and structurally complex, have been isolated from the seed oil, bark and leaves of neem (Daniel and Smith 1991) <sup>[13]</sup>. Phytochemicals are often distasteful and toxic to various insect pests. They can modify feeding deterrence behaviour of an insect (Schmidt and Strelake, 1994) <sup>[14]</sup>. Moreover, as most aphid species have become resistant to many aphidicidal agents (Rashid *et al.* 2013) <sup>[15]</sup> managing these pests in greenhouses and in the field is becoming problematic (Srivastava and Awasthi 1958, Dubey *et al.*, 2004, Dang *et al.* 2010, Kim *et al.* 2011) <sup>[16, 17, 18, 19]</sup>. Therefore, in the present investigation the selected naturally occurring asteraceous plant extractives are efficient and environment friendly pest control alternatives must be developed to replace toxic and hazardous synthetic pesticides.

## 2. Materials and Methods

**(A) Procurement of Raw Plant Materials:** The plant parts used for extraction were surveyed, identified and collected mainly from wasteland, wild areas and some plants were collected from cultivated fields of the farmers. The investigations on the screening of various available indigenous naturally occurring ten plant extracts *viz.*, aerial parts of *Acemella paniculata* Well ex DC, *Cichorium intybus* (L.), *Chromolaena odorata* Linn., *Chrysanthemum cinerariifolium* (trev.) Vis., *Inula racemosa* Hook. F., *Mantisalca duriaeri* Birq. Et Cavill., *Rechardia tingitana* (L.) Roth, *Rhaponticum acaule* (L.) DC, *Scorzonera undulate* Vahl, and *Tagetes minuta* Linn. were screened for their phagorepellent biopotency against nymph and adults of *Aphis gossypii* Glover in laboratory. For carrying out the present studies only ten selected asteraceous botanical Soxhlet extractives were assessed for their phagorepellent Bioefficacy under laboratory conditions.

**(B) Preparation of Powder and their Extraction:** Fresh collected ten green asteraceous plant parts (leaves, Flowers and roots, aerial parts etc) were washed with distilled water and kept in the laboratory for 7 days for shadow air drying before making powder. Electric grinder was used to have coarse powder then these were passed through a 60-mesh sieve to get fine powder. Powders were kept in polythene bags at room temperature and properly sealed to prevent quality loss. For the extraction Soxhlet Apparatus was used; about 20g powder from each category of powder were extracted

with 300 ml of solvents ie. Alcohol and distilled water. Extractions of each category powder were done in about 12 hrs. After soxhlet extraction, the material was run on rotary evaporator (Trivedi and Chandel, 2009) <sup>[40]</sup>. The extracts were concentrated on rotary evaporator by removing the excess solvent under vacuum. After evaporation of solvent with rotary evaporator the remaining extracted material was kept on water bath for removing remaining solvent from the extracts. The extracts were stored at 4 °C prior to application.

- (C) Preparation of Stock Solution and their Insecticidal Formulations:** For stock solution, 50ml. extract in each case was taken into reagent bottles and 50ml. benzene was added in it to dissolve the contents of the botanical materials. The mouth of the bottles were stopper with airtight corks after which, these bottles containing the solutions were kept in refrigerator. Five concentrations (0.25, 0.5, 1.0, 1.5 and 2.0 per cent) were used for experiments on repellent tests in the laboratory conditions. The different concentrations of the herbal extracts were prepared from the stock solution using benzene as solvent and Triton X-100 as emulsifier. The level of solvent and emulsifier were kept constant.
- (D) Apparatus used for Experiment:** Small plastic jars (capacity 50 ml) were used for the experiment, there was one set of two jars joined by clear plastic pipe of 1.00 cm diameter at an angle of 180 degree for each replication. One jar of each set was provided with 10 g of grains given the name 'A' while the other jar was kept empty and given the name 'B'. In jar 'A', the grains treated with extracts were placed, while the jar B remained empty. The jars used for experiment were disinfected with alcohol.
- (E) Experimental Protocol:** The repellent test was carried out under laboratory condition against nymph and adults of *A. gossypii*. The okra leaves and fruites were used as food for the nymph and adults of *A. gossypii*. Leaves were treated with different concentrations (0.25, 0.5, 1.0, 1.5 and 2.0 per cent) for two minutes. The treated leaves were left under electric fan for about half an hour, to make a dry film of the extracts on the leaves for each set of extract and one control. The treated foods were kept in jar (23cm x 10cm) on moist filter paper. The untreated leaves were dipped in Benzene + emulsified water only. Thirty starved nymphs and adults were released in each jar along with control. Three replicates per treatments were maintained. The treated jars either repelled the insects or forced them to move from treated jars (T) to an empty jar (C) through the plastic pipe. The ones found in plastic pipe were considered repelled individuals. Number of nymphs and adults of *A. gossypii*, reached to the food or repelled to untreated empty jar were recorded.

**Table 1:** List of selected asteraceous plants materials used for extraction

Sr. No.	Scientific Name	Vernacular name	Family	Plant parts used
1.	<i>Acmella paniculata</i> (Wall ex DC.) R.K. Jansen	Toothache Plant	Asteraceae	Aerial parts
2.	<i>Chrysanthemum cinerariifolium</i> (trev.) Vis.	Daisy	Asteraceae	Leaves
3.	<i>Chromolaena odorata</i> Linn.	Siam weed	Asteraceae	Leaves
4.	<i>Cichorium intybus</i> (L.)	Chicory	Asteraceae	Roots
5.	<i>Inula racemosa</i> Hook. f	Puskarmul	Asteraceae	Aerial parts
6.	<i>Mantisalca duriaeri</i> Birq. Et Cavill.	Spach	Asteraceae	Roots
7.	<i>Reichardia tingitana</i> (L.) Roth	False sowthistle	Asteraceae	Flowers
8.	<i>Rhaponticum acaule</i> (L.) DC.	Coffee plum	Asteraceae	Flowers
9.	<i>Scorzonera undulate</i> Vahl	Black Salsify	Asteraceae	Flowers
10.	<i>Tagetes minuta</i> Linn.	Wild Marigold	Asteraceae	Leaves

**Table 2:** Formulations of Extracts

Concentration (%)	Amount of Stock Solution (ml)	Amount of Benzene (ml)	Amount of Emulsifiable Water (ml)	Total Amount (ml)
0.25	2.50	22.50	475.00	500.00
0.50	5.00	20.00	475.00	500.00
1.00	10.00	15.00	475.00	500.00
1.50	15.00	10.00	475.00	500.00
2.00	20.00	5.00	475.00	500.00

**Repellent Bioassay:** For testing the Phago-repellent effect of asteraceous extracts, okra leaves were used as food for nymph and adults of *Aphis gossypii* treated with 0.25, 0.5, 1.0, 1.5 and 2.0 per cent concentrations. The treated okra leaves and fruits were kept in jar (23cm x 10 cm) on moist filter paper. Thirty (15 male and 15 female), 24 hours starved nymph and adults of *Aphis gossypii* were released in each jar along with control. For control treated foods were dipped in Benzene + emulsified water only. After four hours of the release nymph and adults of *Aphis gossypii*. Treated nymph and adults of *Aphis gossypii* either repelled and forced them to move from treated jars A to an empty jar B through the plastic pipe The ones found in plastic pipe were considered repelled individuals. The repellency data (in treated and untreated jars) and alive (in empty or untreated jars) were recorded for 6hr at an interval of 6 hours for each observation. The data was collected and calculated on the number of nymph and adults of *Aphis gossypii*, which reached the treated food and repellency over control was recorded and calculated (Finney,

1925)<sup>[20]</sup>.

### 3. Result and Discussion

Data depicted from the results based on actual EC<sub>50</sub> values, the sequence of repellency of different plant extractives against nymph and adults of *A. gossypii* are as: *C. cinerariifolium* (0.0855) > *C. intybus* (0.1359) > *I. racemosa* (0.1531) > *T. minuta* (0.1833) > *M. duriaeri* (0.3208) > *S. undulate* (0.4123) > *A. paniculata*, (0.6365) > *R. acaule* (0.6469) > *R. tingitana* (0.7124) > *C. odorata* (0.9025), respectively., whereas *C. odorata* is taken as a unit. On the other hand data based on relative EC<sub>50</sub> values in the present phago-repellent test against nymph and adults of *A. gossypii*. The sequence of repellency from different plant extractives are as: *C. cinerariifolium* (10.550) > *C. intybus* (6.640) > *I. racemosa* (5.894) > *T. minuta* (4.965) > *M. duriaeri* (2.821) > *S. undulate* (2.188) > *A. paniculata*, (1.417) > *R. acaule* (1.395) > *R. tingitana* (1.267) > *C. odorata* (1.000), times as repellents, whereas *C. odorata* is taken as a unit.

**Table 3:** Calculation of log conc./Probit repellency Regression column of asteraceous extracts on *Aphis gossypii* Glover

Plant Extracts	Heterogeneous	X <sup>2</sup>	Regression equation	EC <sub>50</sub>	Fiducial Limit	Relative EC <sub>50</sub>
<i>Acmella paniculata</i>	3	1.803	0.89X + 0.05	0.6365	M <sub>1</sub> =2.8263 M <sub>2</sub> =1.2632	1.417
<i>Chromolaena odorata</i>	3	1.492	1.08X + 2.65	0.9025	M <sub>1</sub> =2.3362 M <sub>2</sub> =1.9781	1.000
<i>C. cinerariifolium</i>	3	2.834	1.05X + 2.80	0.0855	M <sub>1</sub> =2.3060 M <sub>2</sub> =1.8719	10.550
<i>Cichorium intybus</i>	3	0.123	1.22X + 2.74	0.1359	M <sub>1</sub> =2.0266 M <sub>2</sub> =1.6557	6.640
<i>Inula racemosa</i>	3	1.93	1.41X + 3.26	0.1531	M <sub>1</sub> =1.4512 M <sub>2</sub> =0.2168	5.894
<i>Mantisalca duriaeri</i>	3	4.601	1.101X + 2.89	0.3208	M <sub>1</sub> =2.1173 M <sub>2</sub> =1.7484	2.821
<i>Reichardia tingitana</i>	3	1.080	1.25X + 2.42	0.7124	M <sub>1</sub> =2.0267 M <sub>2</sub> =1.9235	1.267
<i>Rhaponticum acaule</i>	3	1.95	0.92X + 4.37	0.6469	M <sub>1</sub> =1.1588 M <sub>2</sub> =0.0148	1.395
<i>Scorzonera undulate</i>	3	0.42	1.532X + 2.54	0.4123	M <sub>1</sub> =0.0002 M <sub>2</sub> =0.0005	2.188
<i>Tagetes minuta</i>	3	4.601	1.10x+2.89	0.1833	M <sub>1</sub> =2.1173 M <sub>2</sub> =1.7484	4.865

In case of X<sup>2</sup> was found non-significant heterogeneous at P=0.05, Y=Probit repellency, X=Log Concentration X 10<sup>-1</sup>, D.F.=Degree of Freedom, E.C.<sub>50</sub>= Concentration Calculated at given 50% repellency




Fig 1: Log conc./Probit repellency Regression column of *Aphis gossypii* Glover




Fig 2: Log conc./ Probit repellency Regression radar based on EC50 of *Aphis gossypii* Glover.




Fig 3: Log conc./ Probit repellency Regression area based on relative EC50 of *Aphis gossypii* Glover.

4. Discussions

Various botanical products and their extractives works as repellent has been reported by several researchers against *C. chinensis* L., *C. maculatus* and *T. castaneum* etc. (Tripathi, et al. 2000, Valsalan and Gokuldas, 2015) [21, 22] tested effect of plant powders and extracts against *C. chinensis* L. attacking black gram. Certain vegetable oils i.e. turmeric, sweat flag, neem oil and Margosan-O extract 800 ug/ cm<sup>2</sup> applied to filter paper in choice chamber test, produced 59.00 and 67.00 per cent repellency against *T. castaneum* (Jilani et al. 1985, 1988 Mendes et al. 1997, Maredia et al.1992, Pavela, 2006, 2011 and Zoubiri and Aoumeur) [25, 26, 27, 28, 29] *Neem* seed oil at different concentration against *Trichogramma chilonis*. *Neem* seed oil at 0.3 per cent showed high oviposition deterrent

results (Ramamurthy et al. 2002) [30]

In the support of present finding the many entomologist worked on their selected botanicals, among them *Cleome gynandra* gave significant repellent efficacy to spider mite (*Tetranychus urticae* Koch). *Ocimum suave* was found to repel the tick *Rhipicephalus appendiculatus* (Esther et al. 1995 and Abubakar et al. 2000) [31, 32]. Aqueous and methanolic plant extracts of *T. minuta*, which exhibited high repellency (IR = 0.04) against repellent effectiveness against the red flour beetle, *Tribolium castaneum* Herbst (Coleoptera Tenebrionidae). The application of these botanicals may be promising in protecting of stored grains against coleopteran pests. (Padin et al. 2013, Liu et al. 2006) [33, 34]. *Alpinia galanga* L. species of termites, *Coptotermes gestroi* Wasmann and *Coptotermes curvignathus* (Holmgren) (Isoptera: Rhinotermitidae). Repellent activity shows that 250 ppm of 1,8-cineol caused 50.00 ± 4.47% repellency for *C. gestroi* (Fauziah Abdullah et al. 2015) [35]. The maximum repellent activity was observed at 500 ppm in methanol extracts of *N. nucifera*, ethyl acetate and methanol extract of *P. nigrum* and methanol extract of *T. ammi* and the mean complete protection time ranged from 30 to 150 min with the different extracts tested. (Chinnaperuma et al. 2012) [36]. The repellent properties of certain asteraceous plant extractives as stored grain pest (Chandel and Singh,2016, 2017) [37, 38] and crop pests were earlier reported by (Singhamony 1984, Won- Sik, et al. 2002, Trivedi and Chandel, 2009, 2010, Manish Kumar et al. 2017) [39, 40, 41, 42, 43].

5. Conclusion

Conclusively, the present investigation revealed that the appear prospects in selected botanicals *C. cinerariifolium*, *C. intybus*, *I. racemosa* and *Tagetes minuta* were registered promising repellency with minimum EC<sub>50</sub> values to the nymph and adults of *Aphis gossypii*, when compared to other extracts. The data depicted in table 1 and figure 1 indicated that result based on their relative EC<sub>50</sub> values the extracts of *C. cinerariifolium* showed highest relative repellency with minimum relative EC<sub>50</sub> Values (19.550) to the treated nymph and adults of *Aphis gossypii* followed by *C. intybus* (6.39), *I. racemosa* (5.88) and *T. minuta* (5.42) times more repellent than *C. odorata* (1.00), which is taken as unit. The above selected plant materials can be use in protection of okra, *Abelmoschus esculentus* crop from *Aphis gossypii* infestations. The *C. cinerariifolium*, *C. intybus*, *I. racemosa* and *Tagetes minuta* extractives can be applied effectively by

farmers as a component of integrated management of insect-pests of solanaceous vegetable and crop in India.

## 6. Acknowledgement

Authors would like thanks to Principal D.B.S.College, Kanpur and Head, Department of Zoology, D.B.S.College, Kanpur for providing the necessary laboratory facilities. Prof. N.D.Pandey, Head, Division of Entomology, C S Azad University of Agriculture and Technology, Kanpur for rendering their support and help for the completion of this work.

## 7. References

1. Radake SG, Undirwade RS. Seasonal abundance and insecticidal control of shoot and fruit borer, *Earias* spp. on okra, *Abelmoschus esculentus* (L.). Indian J Entomol. 1981; 43:283-287.
2. Adeboye OC, Oputa CO. Effects of galex on growth and fruit nutrient composition of okra (*Abelmoschus esculentus* L. Moench) Int. J Agric. 1996; 18(1-2):1-9.
3. Kumar DS, Tony DE, Kumar AP, Kumar KA, Rao DBS, Nadendla RA. Review On: *Abelmoschus esculentus* (Okra), International Research Journal Pharmaceutical Applied Science. 2013; 3(4):129-132.
4. Konar A, Rai L. Efficacy of some insecticides against shoot and fruit borer *Earias vittella* Fab. and *Earias insulana* (Boisd.) of okra (*Abelmoschus esculentus* L. Moench). Environ. Ecol. 1990; 8(1):410-413.
5. Chaudhary HR, Dadheech LN. Incidence of insects attacking okra and the avoidable losses caused by them. Ann. Arid Zone. 1989; 28(3-4):305-307.
6. Chandel BS, Sudhir Tripathi, Arti Singh. Laboratory bioassay of N- (P-methoxy Phenyl-2 hydroxy salicylimide and N- (p-chlorophenyle) -2 hydroxysalicylimide as repellent against pulse beetle, *Callosobruchus chinensis* L. (Coleoptera: Bruchidae). International Journal of Entomology Research. 2018; 3(1):161-266.
7. Park HJ, Baek MY, Cho JG, Seo KH, Lee GY, Moon SJ. Insecticidal alkaloids on aphids from *Corydalis turtchaninovii* tubers. J Korean Soc. Appl. Biol. Chem. 2011; 54:345-352.
8. Chandel BS Sudhir, Tripathi Arti Singh. Laboratory bioassay of N- (P-methoxy Phenyl-2 hydroxy salicylimide and N- (p-chlorophenyle) -2 hydroxysalicylimide as repellent against pulse beetle, *Callosobruchus chinensis* L. (Coleoptera: Bruchidae). International Journal of Entomology Research. 2018; 3(1):265-270.
9. Bajpai R, Chandel BS. Evaluation of *Adhatoda vasica*, *Acorus calamus* and *Vitex nugendo* as a source of phago-antifeedant against cabbage butterfly, *Pieris brassicae* Linn. (Lepidoptera: Pieridae) on mustard, *Brassic campestris* (Linn.). International Journal of Biotechnology and Biochemistry, 2009; 5:451-459.
10. Bajpai R, Chandel BS. Assessment of certain plant extracts as insecticides against cabbage butterfly, *Pieris brassicae* Linn. (Lepidoptera: Pieridae). International Journal of Biotechnology and Biochemistry, 2010; 6(6):1003-1017
11. Antonius AB, Hagazy G. "Feeding deterrant activity of certain botanical extracts against cotton leafworm, *Spodoptera litralis* Baised." Am. Agric. Sci. 1987; 32(1):719-729.
12. Chandel BS, Sengar HS. Insectifuge effect of neem, *Azadirachta indica* A. Juss and their derivatives against sawfly, *Athalia lugens proxima* Klug. (Hymenoptera: Tenthredinidae). International Journal of Zoology Studies. 2018; 3(1):53-59.
13. Daniel SH, Smith RH. The repellent effect of neem (*Azadirachta indica* A. Juss) oil and its residual efficacy against *Callosobruchus maculatus* (Coleoptera: Bruchidae) on cowpea. Fleurat-Lessard, F. and Ducom, P., Eds. Proceedings Fifth International Working Conference on Stored-Product Protection, 1991, 1589-1598.
14. Schmidt GH, Streloke M. Effect of *Acorus calamus* Linn. (Araceae) oil and its main compound B- Asarone on *Prostephanus truncates* Horn (Coleoptera: Bostrichidae) J Stored Product Research. 1994; 30(3):227-235.
15. Rashid T, Chen J, Mcledo P. Toxicity of newly isolated piperidine alkaloids from the red imported fire ant, *Solenopsis invicta*, against the green peach aphid, *Myzus persicae* (Sulzer). Adv. Entomol. 2013; 1:20-23
16. Srivastava AS, Awasthi GP. An insecticide from the extract of the plant *Adhatoda vesica* harmless to man." P r o c. 10<sup>th</sup> Ent. Congr. Expi. 1958; 2:245-246.
17. Dubey A, Gupta R, Chandel, BS. Efficacy of *Acorus calamus*, *Vitex negundo* and *Ageratum conyzoides* against tobacco caterpillar *Spilarctia obliqua* walker." Indian. J Entomology. 2004; 66(3):238-240
18. Dang QL, Lee GY, Choi YH, Choi GJ, Jang KS, Park M. et al. Insecticidal activities of crude extracts and phospholipids from *Chenopodium ficifolium* against melon and cotton aphid, *Aphis gossypii*. Crop Prot. 2010; 29:1124-1129.
19. Kim SK, Kim YC, Lee S, Kim JC, Yun MY, Kim IS et al. Insecticidal activity of rhamnolipid isolated from pseudomonas sp. EP-3 against green peach aphid, J Agric Food Chem. 2011; 9, 59(3):934-938
20. Finney DJ. "Probit analysis: a statistical treatment of the sigmoid responses in curve." Rev. ed. Cambridge University Press, London, 1938, 318.
21. Tripathi AK, Prajapati V, Agarwal KK, Khanuja SPS, Kumar S. Repellency and toxicity of oil from *Artemisia annua* to certain stored product beetles. Jour Econ. Ent. 2000; 93(1):43-47.
22. Valsalan KK, Gokuldas M. Repellent and oviposition deterrent effects of *Clerodendrum infortunatum* on the pulse beetle *Callosobruchus chinensis* L. (Coleoptera: Bruchidae), Journal of Entomology and Zoology Studies. 2015; 3(4):250-255.
23. Jilani G, Saxena RC, Rueda BP. Repellency and toxicity of some plant oils and their terpene components to *Sitotroga cerealella* Oliver (Lipidoptera, Gelechiidae), Tropical Science. 1985; 25:249-252.
24. Jilani G, Saxena RC, Rueda BP. Repellent and growth inhibiting effects of turmeric oil, sweetflag oil, *Neem* oil and Margosan oil on red flour beetle (Coleoptera: Tenebrionidae). Jour Econ. Ent. 1988; 81(4):1226-1230.
25. Mendes RO Queiroz, Zani CL. Screening of asteraceae (Compositae) plant extracts for larvicidal activity against *Aedes fluviatilis* (Diptera: Culicidae). Mem. Inst. Oswaldo Cruz, 1997; 92:565-570.
26. Maredia KM, Segura OL, Mihm JA. Effects of neem,

- Azadirachata indica*, on six species of maize insect pests. Tropical Pest Management. 1992; 38(2):190-195.
27. Pavela R. Insecticidal activity of essential oils against cabbage aphid *Brevicoryne brassicae*. J Essent. Oil Bearing Plant. 2006; 9:99-106.
  28. Pavela R. Insecticidal and repellent activity of selected essential oils against of the pollen beetle, *Meligethes aeneus* (Fabricius) adults. Ind. Crop. Prod. 2011; 34:888-892
  29. Zoubiri Safia, Baaliouamer Aoumeur. Potentiality of plants as source of insecticide principles Journal of Saudi Chemical Society. 2015; 19(1):1-116.
  30. Ramamurthy R Rajaram V, Rajendran LM. Efficacy of plant products against the storage pest bruchides in black gram. Legume Research, 2002, 32:34.
  31. Esther N Mwangi, Ahmed Hassanali, Suliman Essuman, Edward Myandat, Lambert Moreka, Mark Kimondo *et al.* Repellent and acaricidal properties of *Ocimum suave* against *Rhipicephalus appendiculatus* ticks. Experimental & Applied Acarology. 1995; 19(1):11-18.
  32. Abubakar MS, Abdurahman EM, Haruna AK. The repellent and antifeedant properties of *Cyperus articulatus* against *T. castaneum*. Phytotherapy Research. 2000; 14:281-283.
  33. Padin SB, Fuse C, Urrutia MI, Dal Bello GM. Toxicity and repellency of nine medicinal plants against *Tribolium castaneum* in stored wheat, Bulletin of Insectology. 2013; 66(1):45-49,
  34. Liu CH, Mishra AK, Tan RX Repellent. Insecticidal and phytotoxic activities of isoalantolactone from *Inula racemosa*. Crop Protection, 2006; 25:508-511.
  35. Fauziah Abdullah, Partiban Subramanian, Halijah Ibrahim, Sri Nurestri, Abdul Malek, Guan Serm Lee *et al.* Chemical composition, antifeedant, repellent, and toxicity activities of the rhizomes of galangal, *Alpinia galanga* against Asian subterranean Termites, *Coptotermes gestroi* and *Coptotermes curvignathus* (Isoptera: Rhinotermitidae). J Insect Sci. 2015; 15(1):322-437
  36. Chinnaperumal Kamaraj, Abdul Abdul Rahuman, Asokan Bagavan, Gandhi Elango, Abdul Abduz Zahir, Thirunavukkarasu *et al.* Larvicidal and repellent activity of medicinal plant extracts from Eastern Ghats of South India against malaria and filariasis vectors. Asian Pacific Journal of Tropical Medicine. 2012; 4(9):698-705
  37. Chandel BS. Arti singh. Repellent biopotency of *Cichorium intybus*, *Inula racemosa*, *Tagetes minuta* and *Mantisalca duriaeri* aquash formulations against pulse beetle, *Callosobruchus chinensis* Linn. (Coleoptera: Bruchidae). Life Science Bulletin. 2016; 13(2):65-68.
  38. Chandel BS, Singh A. Phagodeterrent bioefficacy of *Acorus calamus*, *Withania somnifera* and *Momordica charantia* against chickpea bruchids, *Callosobruchus chinensis* (Linn.) (Coleoptera: Bruchidae), Journal of Entomology and Zoology Studies. 2017; 5(6):935-939.
  39. Sighamony S, Anees I, Chandrakala TS. Osmani Z. Natural products as repellents for *Tribolium castaneum* Herbst. International Pest Control, 1984, 156-157
  40. Won-Sik C, Park BS, Ku SK, Lee SE. Repellent activities of essential oils and monoterpenes against *Culex pipiens pallens*. J Am. Mosquito Control Assoc. 2002; 18:348-351.
  41. Trivedi SS, Chandel BS. Repellent Efficacy of *Adhatoda vasica*, *Pongamia glabra* and *Momordica charantia* plant extract against *Spilarctia obliqua* Walker (Lepidoptera: Arctiidae). International Journal of Biotechnology and Biochemistry. 2009; 4(4):469-461.
  42. Trivedi SS, Chandel BS. Antifeedant biopotency of certain indigenous naturally occurring plant extractives against Bihar hairy caterpillar, *Spilarctia obliqua* Walk. (Lepidoptera: Arctiidae). International Journal of Biotechnology and Biochemistry. 2010; 6(6):1019-1026.
  43. Kumar M, Tripathi RC, Chandel BS. Potential impact of *Inula racemosa*, *Cichorium intybus* and *Mantisalca duriaeri* against *Plutella xylostella* Linn. (Lepidoptera: Noctuidae). Life Science Bulletin. 2017; 14(1):65-69.