



# International Journal of Fauna and Biological Studies

Available online at [www.faunajournal.com](http://www.faunajournal.com)

I  
J  
F  
B  
S

International  
Journal of  
Fauna And  
Biological  
Studies

ISSN 2347-2677

[www.faunajournal.com](http://www.faunajournal.com)

IJFBS 2020; 7(4): 115-119

Received: 15-05-2020

Accepted: 19-06-2020

**Masroof Ahmad Malik**

Ph.D., Research Scholar,  
Department of Bio-Chemistry  
J. S. University Shikohabad,  
Utter Pradesh, India

**Altaf Ahmad Ganaie**

Ph.D., Research Scholar,  
Department of Bio-Chemistry  
J. S. University Shikohabad,  
Utter Pradesh, India

## An epidemiological study of gastrointestinal protozoan parasites among children

**Masroof Ahmad Malik and Altaf Ahmad Ganaie**

### Abstract

The study was intended to explore the prevalence of Gastrointestinal Protozoan Parasites (PPP) among children with special reference to their source and type of drinking water. 400 respondents were selected with the help of convenient sampling technique. The study was carried in context of descriptive research method. 400 respondents were selected from Kulgam Districts of Union Territory of Jammu and Kashmir. Further, all respondents were selected within the age group of 12-16 years. The results of the study reveal that that type of drinking water has significant impact on the prevalence of Gastrointestinal Protozoan Parasites (GPP) among children. Besides, it was found that source of drinking water has significant impact on the prevalence of Gastrointestinal Protozoan Parasites (GPP) among children. The impact of defecation site was reported significant on the prevalence of Gastrointestinal Protozoan Parasites (GPP).

**Keywords:** Gastrointestinal Protozoan Parasites (GPP) Source of Drinking Water, Type Drinking Water, Defecation Sites

### 1. Introduction

Epidemically speaking, intestinal parasitic diseases are worldwide in dissemination and especially endemic in nations which are still coming under developing nations list. Very high pervasiveness in developing nations is because of absence of sanitation, absence of hygiene therefore, such kind of epidemiology occurs wherever there is unhygienic environment, open defecation site, lack of health consciousness etc. Individuals of any age are influenced by this pattern of predominant parasitic diseases; notwithstanding, children are the most exceedingly terrible. It is the most pervasive protozoan parasite influencing around 200 million individuals, as of now contaminated. Affected individuals are 3.5 billion, and people that are sick, their number is about 450 million are sick because of these contaminations. These contaminations are observed as a severe public health problem, as they cause delayed and stunted development in children and health problems. There are large numbers of factors associated with rapid increase of gastrointestinal parasites infection in children. There is direct relationship between malnutrition and intestinal parasitism and between nutritional status and both constancy and potency of infections. Malnutrition may affect many developmental processes in children such as children development, cognitive function is also affected and thus affecting school performance of children. It causes complications such as, severe anaemia, intestinal blockage etc. The prevalence of malnutrition, lack of awareness, unhygienic environment and degraded eating habits are some of the factors which are seen in India for facilitating the occurrence of gastrointestinal parasites infection especially in children. Keeping in view, the apex institution of health, World Health Organization (WHO) stated that amoebic infection causes around 50 million people in the world to suffer each year, which results in 40100 thousand deaths yearly. Parasitic infection is more common in developing countries including India. Some parasites don't noticeably affect their host while some others become mature, reproduce or attack organ system which makes their host ill, resulting into a parasitic infection. Although all infectious agents in human are considered as parasites but in convention, parasitic diseases are defined as those caused by protozoa or helminthes. Lack of adequate medical care facilities and social marginalization has increased the susceptibility of the population to other pathogenicities and morbidities associated with parasitic infection<sup>[5]</sup>. Parasites can live throughout the body but mostly prefer the intestinal wall. Apart from causing morbidity and mortality, intestinal parasites infection have been associated with stunting, physical weakness, open defecation sites, impure water, low educational performance of school children.

**Corresponding Author:**

**Masroof Ahmad Malik**

Ph.D., Research Scholar,  
Department of Bio-Chemistry  
J. S. University Shikohabad,  
Utter Pradesh, India

Intestinal parasite has been predicted to affect some 3.5 billion people worldwide and 450 million are thought to be sick as a result of such infections, the majority is children. In some tropical areas, the prevalence hikes nearly to 100 percent. It is a main socio-economic problem in India, though a hospital based study has shown a falling trend during a decade. Intestinal parasites even in low or moderate number effect on both nutritional and thereby on immune system of individuals leading to numerous morbidity and mortality. Indeed large number of the research studies has been conducted by the researchers on parasites. Like the studies conducted by Al-Saeed, A. T. And Issa, S.H. (2006)<sup>[1]</sup>, Alver, O. And Tore, O. (2006)<sup>[2]</sup>, Ayalew A, Debebe T, (2012)<sup>[9]</sup>, Ayalew *et al.* (2011)<sup>[10]</sup>, Escobedo, A.A.; Canete, T. & Nunez, F.A. (2008)<sup>[19]</sup>, Escobedo, A. J. (2010)<sup>[20]</sup>, Hegazy, A. M., Younis, N.T.; & Badr, A.M. (2014)<sup>[21]</sup>, Jain M, Jain J, Gupta S. (2016)<sup>[23]</sup> and Jarallah, H.M. (2012)<sup>[24]</sup>. However, there may be hardly nay study which will be in the study area of Kulgam District. In context to same, the investigator explored the below mentioned research problem.

**1.2 Research Problem:** The statement of the research problem is as under:

“An Epidemiological Study of Gastrointestinal Protozoan Parasites among Children”

**1.3 Objectives of the study:** The objectives of the present study are as under:

- **Objective I:** To explore the prevalence of Gastrointestinal Protozoan Parasites among Children With Special Reference To Their Source of Drinking Water.
- **Objective II:** To explore the prevalence of Gastrointestinal Protozoan Parasites among Children With Special Reference To Their type of Water usage.
- **Objective II:** To explore the prevalence of Gastrointestinal protozoan parasitic infection in relation to defecation Site.

**1.4 Research question:** The aim for the presents study was to explore the impact Source and Type of Drinking Water on prevalence Gastrointestinal Protozoan Parasites among children. Besides, it aims the impact of defecation sites on prevalence Gastrointestinal Protozoan Parasites (GPP) among children. In pursuance to same, it was presumed that the selected variables has significant impact on prevalence Gastrointestinal Protozoan Parasites (GPP) among children.

**1.5 Operational definition of terms and variables:** The operational definitions of terms and variables are as under:

1. **Gastrointestinal Protozoan Parasites:** In the present study gastrointestinal protozoan parasites refers dominant set of scores obtained by respondents on stool sample tests and blood test. Besides, the investigator employed a self-developed questionnaire (2020).
2. **Source of water:** Source of drinking water in the present study refers the water availed by respondents like pond water, river water, stream water and well water.
3. **Type of water:** Type of drinking water I the resents study refers the dichotomy made on the basis of boiled an uncoiled water.
4. **Defecation Site:** Defecation Site in the presents study refers the defecation Site availed by the respondents like open field, pit Latrines, sanitary latrines.

**1.6 Research limitations of the study:** The present study will be delimited two 400 respondents of Kulgam districts. Besides, whole sample was sleeted from the above mentioned sampling sites of Kulgam district of Union Territory of Jammu and Kashmir. further, all respondents were elected within the age group of 12-16 years.


**1.7 Rationale of the study:** Keeping the feasibility and usability of the study under consideration, the researcher found it suitable to go through descriptive survey method. Accordingly, present study was carried with the help of descriptive method. The parameters involved in methodology and procedure are as under:

- **1.7.1 Sample:** 400 respondents were selected with due representation of gender of school. The age groups of the subjects were ranged 18-21 years. Whole data was selected for the present study.
- **1.7.2 Sampling technique:** Whole data was selected by using Convenient Sampling Technique (CST). The required data was collected from District Kulgam. The below mentioned table indicates the precise explanation of sample:

**Table 1:** Showing the selection of sample with dichotomy representation

Sampling Site	Yaripora	Frisal	Devsar	Quimoh	Pahloo	DK Pora
Sample	90	40	60	80	60	70
Total= 400						

- **1.7.3 Measuring Instruments:** All the three components were assessed separately the detailed analyses of these instruments is reported as under:




**Fig 1:** Showing the selection of sample with dichotomy representation

**1.8 Analysis and interpretation of the data:** In connection to same, collected data was put to suitable statistical treatment by using Mean, S. D. and ‘t’ value. The detailed procedure of statistical treatment is analysed as under: The analysis and interpretation of the data is reported as under:

**Table 2:** Prevalence of Gastrointestinal protozoan parasitic infection in relation to Source of Drinking Water. (N=400)

Variable	Tap water		Well water		River/Pond/ Stream	
	Examined	Infected	Examined	Infected	Examined	Infected
source of Water	300	30	70	10	30	50
	Frequency	Percentage	Frequency	Percentage	Frequency	Percentage
	30.00	7.5	70.00	2.5	30.00	12.5




**Fig 2:** Showing graphical representation of prevalence of gastrointestinal protozoan parasitic infection in relation to Source of Drinking Water. (N=400)

**Interpretation 1.2:** While glancing towards table 1.2, (Please consult Table 2, Fig. 2) gives information about the frequency and percent wise distribution of gastrointestinal protozoan parasite infection in relation to their source of water. The results reveal that those respondents who use tap water 7.5% (F=30) were seen with gastrointestinal protozoan parasite infection. According the results reported in the same table reveals that among 70 respondents were found who avail well water in their day to day life schedule and among these respondents 10% (F=70) were seen with gastrointestinal protozoan parasite infection. The obtained results indicate that among river/pond or stream water users 12.5% (F=30) were seen with gastrointestinal protozoan parasite infection. Thus, from the obtained results the investigator can generalize that river/pond and stream water users were found with high gastrointestinal protozoan parasite infection rate. Next to it tap water users were found with gastrointestinal protozoan parasite infection rate. Thus, type of water resources were reported to have a major impact on the level of gastrointestinal protozoan parasite infection. The results may attribute to this fact that open water sources like rivers, ponds and streams are more prone to parasitic contamination.

**Table 3:** Prevalence of Gastrointestinal protozoan parasitic infection in relation to Water usage. (N=400)

Variable	Boiled water			Un-boiled water		
	Examined	Infected	Percentage	Examined	Infected	Percentage
Water Usage	40	2	0.5	360	88	22




**Fig 3:** Showing graphical representation of Gastrointestinal protozoan parasitic infection in relation to water usage.

**Interpretation:** While glancing towards table 1.3, (Please consult Table 3, Fig. 3) gives information about the frequency and percent wise distribution of gastrointestinal protozoan parasite infection in relation to their water usage. The results reveal that among those respondents who avail boiled water, 0.5% (F=40) were seen with gastrointestinal protozoan parasite infection. While analysing those respondents who avail un-boiled water, it was found that 22% (F=88) were seen with gastrointestinal protozoan parasite infection. Thus, from the reported results, it can be inferred that type of water usage of the respondents hold high level of impact on the level of gastrointestinal protozoan parasite infection of the respondents. As reported from the above results it is obvious that boiled water reduce the concentration of the gastrointestinal protozoan parasite infection among children.

**Table 4:** Showing Prevalence of Gastrointestinal protozoan parasitic infection in relation to defecation Site (N=400)

Defecation Site	Open fields		Pit Latrines		Sanitary Latrines	
	Examined	Infected	Examined	Infected	Examined	Infected
	Frequency	Percentage	Frequency	Percentage	Frequency	Percentage
Site	15	3.75	50	12.5	25	6.25



**Fig 4:** Showing graphical representation of Prevalence of Gastrointestinal protozoan parasitic infection in relation to Defecation Site

**Interpretation:** The fleeting look on the table 4: (Please Consult table 4) gives information about the Prevalence of Gastrointestinal protozoan parasitic infection in relation to defecation site availed by the respondents. The results reveal that 20 respondents were seen who avail defecating in open fields. However, among these respondents 3.75% (F=15) were found with Gastrointestinal protozoan parasitic infection. In context to same, it was found that 150 respondents were seen who were defecating in pit latrines. Hover, among these respondents 12.5% (F=50) were seen with Gastrointestinal protozoan parasitic infection. Besides the calculated results indicate that 230 respondents were seen who avail sanitary latrines, but 6.25% (F= 25) were reported with Gastrointestinal protozoan parasitic infection. Thus, from the gained results it can be inferred that open field defecation and pit latrines defecation were seen more prone to Gastrointestinal protozoan parasitic infection as compared to sanitary latrines.

**2. Conclusion:** The aim for the presents study was to explore the impact Source and Type of Drinking Water on prevalence Gastrointestinal Protozoan Parasites among children. Besides, it aims the impact of defecation sites on prevalence Gastrointestinal Protozoan Parasites (GPP) among children.

In pursuance to same, it was found that type of drinking water has significant impact on the prevalence of Gastrointestinal Protozoan Parasites (GPP) among children. Besides, it was found that source of drinking water has significant impact on the prevalence of Gastrointestinal Protozoan Parasites (GPP) among children. Further, it was found that the impact of defecation site has significant impact on the prevalence of Gastrointestinal Protozoan Parasites (GPP) among children.

- **Support to findings:** Indeed, it was found that type of drinking water has significant impact on the prevalence of Gastrointestinal Protozoan Parasites (GPP) among children. Besides, it was found that source of drinking water has significant impact on the prevalence of Gastrointestinal Protozoan Parasites (GPP) among children. The results are supported by host of the researchers like; Al-Saeed, A. T. And Issa, S.H. (2006)<sup>[1]</sup>, Alver, O. And Tore, O. (2006)<sup>[2]</sup>, Ayalew A, Debebe T, (2012)<sup>[9]</sup>, Ayalew *et al.* (2011)<sup>[10]</sup>, Escobedo, A.A.; Canete, T. & Nunez, F.A. (2008)<sup>[19]</sup>, Escobedo, A. J. (2010)<sup>[20]</sup>, Hegazy, A. M., Younis, N.T.; & Badr, A.M. (2014)<sup>[21]</sup>, Jain M, Jain J, Gupta S. (2016)<sup>[23]</sup> and Jarallah, H.M. (2012)<sup>[24]</sup>.
- **Conflict of Interest:** Keeping the entire research process under consideration, no any conflict of interest has been declared by the investigator.

### 3. References

1. Al-Saeed AT, Issa SH. Frequency of Giardia Lamblia Among Children in Doihuk, Northern Iraq. Eastern Mediterranean Health Journal. 2006; 12(5):555-562.
2. Alver O, Tore O. The Prevalence and Distribution of Intestinal Parasites Detected by The Uludag University Medical School. *Turkiye Parazitol Derg.* 2006; 30(4):296-301.
3. Alzain BF, Sharma PN. Hemoglobin Levels & Protozoan Parasitic Infection in School Children of Udaipur City. *Journal of Al Azhar University-Gaza (Natural Sciences). Journal of Medical Laboratory Sciences*, 2(2):74-76.
4. Anvarian M. The Prevalence of Intestinal Protoza in Partients Referred to Tabriz Imam Reza Hospital from January 2010 To December 2010. *Advances In Environmental Biology.* 2011; 5(&):1916-1919.
5. Ashatiani *et al.* Prevalence of Intestinal Parasites Among Children Referred to Childrens Medical Centre During 18 Years (1991-2008), Tehran, Iran *Ann Trop Med Parasitol.* 2011; (7):507-513.
6. Assudani H, Gusani J, Agravat H. Intestinal Parasitic Infections in Paediatric Patients with Diarrhea with Special Emphasis to Opportunistic Parasites and Predisposing Factors. *Journal of Al Azhar University-Gaza.* 2015; 8(10):35-40.
7. Attias *et al.* Current Caregiver Awareness of Pediatric Giardiasis and Cryptosporidiosis. *Journal of Al Azhar University-Gaza.* 2015; 8(10):35-40.
8. Awasthi S, Pand VK. Prevalence of Malnutrition and Intestinal Parasites in Preschool Slum Children in Lucknow. *Indian Pediatr. Journal of Al Azhar University-Gaza.* 1997; 8(10):35-40.
9. Ayalew A, Debebe T, Worku A. Prevalence and Risk Factors of Intestinal Parasites Among Delgi School Children, North Gondar, Ethiopia. *Journal of Parasitology and Vector Biology.* 2012; 3(5):75-81
10. Ayalew *et al.* Prevalence and Risk Factors of Intestinal Parasites Among Delgi School Children, North Gondar, Ethiopia. *Journal of Parasitology and Vector Biology.* 2011; 3(5):75-81
11. Bahadur Sarkari *et al.* Prevalence and Risk Factors of Intestinal Protozoan Infections. A Population -Based Study in Rural Areas of Boyer- Ahmad District, Southwestern Iran *BMC Infections Disesaes. Journal of Al Azhar University-Gaza.* 2016; 8(10):35-40.
12. Baldo ET, Belizario VY, Leon WUD, Kong H, Chung D. Infection Status of Intestinal Parasites In Children Living In Residential Institutions In Metro Manila, The Philippines. *The Korean Journal of Parasitology.* 2016; 3(5):75-81.
13. Baldo ET, Belizario, Chung D. Infection Status of Intestinal Parasites in Children Living in Residential Institutions in Metro Manila, The Philippines. *The Korean Journal of Parasitology.* 2004; 42(2):67-70.
14. Elvia Diaz, Rosamaria Bernal. Epidemiology & Control of Intestinal Parasites with Nitazoxanide in Children in Mexico. *The American Journal of Tropical Medicine & Hygine.* 2003; 68(4):384-385.
15. Emile N, Bosco NJ, Karine B. Prevalence of Intestinal Parasitic Infections & Associated Risk Factors Among Kigali Institute of Education Students in Kigali, Rwanda *Tropical Biomedicine.* 2013; 30(4):718-726.
16. Ensink JHJ, Hoekc WV, Amerasinghec FP. Giardia Duodenalis Infection & Wastewater Irrigation in Pakistan. *Royal Society of Tropical Medicine & Hygiene.* 2006; 30(4):718-726.
17. Eric Buteraet Al. Prevalence of Risk Factors of Intestinal Parasites Among Children Under Two Years of Age in A Rural Area of Rutsiro distinct, Rwanda. *The Pan African Medical Journal.* 2019; 30(4):718-726.
18. Ertan P, Yereli K, Kurt O, Balcioglu C, Ali Onag A. Serological Levels of Zinc, Copper & Iron Elements Among Girdia Lambilia Infected Children In Turkey. *Pediatrics International.* 2002; 30(4):718-726.
19. Escobedo AA, Canete T, Nunez FA. Prevalence, Risk Factors & Clinical Features Associated With Intestinal Parasitic Infections In Children From San Juan Y Martinez, Pinar Del Rio, Cuba. 2008; 30(4):718-726.
20. Escobedo AJ. Giardiasis: The Ever Present Threat of a Neglected Disease. *Infectious Disorders & Drug Targets.* 2010; 10(4):818-826.
21. Hegazy AM, Younis NT, Badr AM. Prevalence of Intestinal Parasites & Its Impact on Nutritional Status Among Pre-School Children Living in Damanhur City,El-Behera Governorate, Egypt. *International Journal of Engineering Science and Innovative Technology (IJESIT).* 2014; 2(3/60):471-485.
22. Hooshyar H, Bagherian T, Baghbani F. Prevalence of Intestinal Parasitic Infections Among Patients Referred to Kashan Reference Laboratory in 2007-2011. *Jundishapur Jhealth Sci.* 2013; 5(1):17-22.
23. Jain M, Jain J, Gupta S. Amoebic Liver Abscess in Children-Experience from Central India. *International Journal of Engineering Science and Innovative Technology (IJESIT).* 2016; 2(3):471-485.
24. Jarallah HM. Intestinal Parasitic Infections Among Rurla Villages in Bashrah Marshes Regions. *Journal of Basrah Researches (Sciences).* 2012; 38(2):40-43
25. Moffat T. Diarrhea, Respiratory Infections, Protozoan Gastrointestinal Parasites & Child Growth in Kathmandu,

- Nepal. The Indian Journal of Pediatrics. 2004; 46(10):358-362.
26. Monajemzadeh SM, Monajemzadeh M. Comparison of Iron & Haematological Indices in Giardia Lamblia Infection Before & After Treatment in 102 Children in Ahwaz, Iran. Med Sci Monit. 2008; 14(1):19-23.
27. Monochitra K, Padukone S, Philips SA, Parija SC. Prevalence of Intestinal Parasites Among Patients Attending A Tertiary Care Center in South India. The Indian Journal of Pediatrics. 2016; 46(10):358-362.