



E-ISSN 2347-2677

P-ISSN 2394-0522

[www.faunajournal.com](http://www.faunajournal.com)

IJFBS 2021; 8(3): 24-27

Received: 06-03-2021

Accepted: 10-04-2021

**Deepak Singh**

Department of Animal  
Husbandry and Dairying,  
C. S. Azad University of  
Agriculture and Technology  
Kanpur, Uttar Pradesh, India

**Ved Prakash**

Department of Animal  
Husbandry and Dairying,  
C. S. Azad University of  
Agriculture and Technology  
Kanpur, Uttar Pradesh, India

**PK Upadhyay**

Department of Animal  
Husbandry and Dairying,  
C. S. Azad University of  
Agriculture and Technology  
Kanpur, Uttar Pradesh, India

**Narendra Kumar**

Department of Animal  
Husbandry and Dairying,  
C. S. Azad University of  
Agriculture and Technology  
Kanpur, Uttar Pradesh, India

**Satendra Kumar**

Department of Animal  
Husbandry and Dairying,  
C. S. Azad University of  
Agriculture and Technology  
Kanpur, Uttar Pradesh, India

**Raj Kumar**

Department of Soil Conservation  
and Water Management,  
C. S. Azad University of  
Agriculture and Technology  
Kanpur, Uttar Pradesh, India

**Corresponding Author:**

**Deepak Singh**

Department of Animal  
Husbandry and Dairying,  
C. S. Azad University of  
Agriculture and Technology  
Kanpur, Uttar Pradesh, India

## Existing breeding management practices by the respondents of Kanpur Nagar, Uttar Pradesh

**Deepak Singh, Ved Prakash, PK Upadhyay, Narendra Kumar, Satendra Kumar and Raj Kumar**

### Abstract

The result of buffalo breeding in the different two blocks Bilhaur and Kalyanpur. These two blocks selected from Kanpur Nagar district of Uttar Pradesh revealed that in the study only 28.00 per cent of the respondents rearing descript buffalo in the study area. It was concluded that in my study more than half 51.00 per cent of the respondents adopted both A.I. and natural breeding and most of the 42.50 per cent respondents were heat detection by mucus discharge + bellowing. The majority of the respondents 52.50 per cent were reported expelled placenta within two hours. It was found according to my village's study noted that buffaloes were allowed for insemination at mid-heat by 82.00 per cent of respondents and pregnancy diagnosis not done by most of the farmers 62.00 per cent in the villages. Most of the respondents 57.50 per cent animals take 3-4.5 year time for first calving. 81.50 per cent of respondents treated the anoestrus as their animals in the villages. Mostly 79.00 per cent of the respondent's buffaloes were taken more than 18-month calving interval. These all information obtained from 200 respondents of the district of Uttar Pradesh.

**Keywords:** Respondents rearing, discharge + bellowing, anoestrus, bubalus bubalis

### Introduction

Buffaloes in the world are categorized into two species: the numerous Asiatic water buffalo (*Bubalus Bubalis*) and African buffalo (*Syncerus caffer*). Asiatic Water buffalo two subspecies are known, the river ( $2n = 50$ ) and the swamp ( $2n = 48$ ) buffalo. River buffalo are the most numerous species include the Murrah, Jafarabady, Surti, Mehsana, Egyptian and Mediterranean breeds (Iannuzzi, 1994) [6]. India is leading the world in milk production since 1998 and produced 165.4 million tonnes of milk with per capita availability of milk 335 grams during 2016-17. Buffalo shared about 49.2 per cent of the total milk produced in India (Anonyms, 2018) [1]. Buffaloes contribute 30 per cent total meat in the country. Buffaloes are great producers of milk and fulfill the great demand of meat. The major states of buffalo meat production are Uttar Pradesh followed by Andhra Pradesh and Maharashtra.

In the encouragement of the national economy of India, the Livestock sector plays a major role and also increases the growth rate. Due to the lack of detailed information on the animals farming existing breeding practices adopted for different categories of livestock, it's not possible for the policy planners to give full attention to these important aspects of dairy cattle production. Efficient management requires a strong database. Efforts should aim to collect information related to animal rearing. Few efforts are known to have been made to study systematic information related to livestock farming and its management adopted in the village. This has not been adequate to serve as the basis on which valid guidelines for introducing scientific managerial practices for improvements of livestock dairy animals can be framed.

### Materials and Methods

The study was conducted purposively in the central zone of Uttar Pradesh. Kanpur occupies the north-western part of the Kanpur division. The present study was conducted by comprising two different blocks Bilhaur and Kalyanpur of Kanpur Nagar district of U.P. The study area has more buffalo population. In each block, Ten villages were selected, and also in all these villages ten farmers were selected randomly have vast opportunities for rural development through buffalo. During summer the temperature of the Kanpur district is very high temperature compared to another district of U.P. The Kanpur lies 132m above sea level and the climate in Kanpur is warm and temperate.

When compared with winter, the summers have much more rainfall. The average annual temperature is 25.6 °C | 78.1 °F in Kanpur. The annual rainfall is 820 mm (32.3 inches). Statistical tools percentage using the methods suggested by Snedecor and Cochran (1994).

## Results and Discussion

The finding regarding various breeding practices followed by respondents in given Table-1.

The finding regarding various breeding practices followed by respondents in given table 1. The role of breeding is very important to generate sufficient stock of improved germplasm for better productivity and healthy animal in the district. The overall result comes from table 1. Indicated that 72.00 per cent were reared non-descript buffaloes only 28.00 per cent reared descript buffaloes like Murra, Bhadawari. This finding is well comparable with the findings of Dhaka *et al.* (2017)<sup>[4]</sup>. That is because descript buffaloes are costly and respondents were not able to pay more money.

It was concluded that in my study more than half 51.00 per cent of the respondents adopted both A.I. and natural breeding, while 36.50 per cent preferred only A.I. and 12.50 per cent respondents adopted natural service in the block of

the district. The present findings were in close agreement with those reported by Sekhar *et al.* (2017)<sup>[12]</sup>, Yadav *et al.* (2016)<sup>[19]</sup>. Some of the contradicted by Chakravarthi *et al.* (2017)<sup>[2]</sup>, Godara *et al.* (2018b)<sup>[5]</sup> found that most of the respondents used A.I. practices at the time of breeding.

Heat detection was observed that by the respondents 42.50, 27.50, 20.00, and 10.00 per cent through mucus discharge + bellowing, mucus discharge, mucus discharge + bellowing + mounting and off feed + urine discharge as the main sign of estrous. This finding is well comparable with finding of Khadda *et al.* (2017)<sup>[7]</sup>, Kumar *et al.* (2019a), Kumar *et al.* (2020)<sup>[8]</sup>, Singh (2018)<sup>[18]</sup>, Viswkarma *et al.* (2018)<sup>[18]</sup>, Yadav *et al.* (2016)<sup>[19]</sup> respondents used mucus and bellowing together for heat detection in animals. Mucus discharge and bellowing is the main sign and respondents assure that buffalo in heat.

The study shows that the majority of the respondents 52.50 per cent was reported expelled placenta within two hours and followed that by the 33.00 per cent take 1 hour after calving but 14.50 per cent take more than 2 hours in the study area. The observation and findings of the study were in favor of finding reported by Singh *et al.* (2016)<sup>[11]</sup>.

**Table 1:** Existing Breeding Management Practices

Existing practices	Bilhaur	Kalyanpur	Respondents	Per cent
<b>1. Breed of buffalo</b>				
A. Descript buffalo	23	33	56	28.00
B. Non-descript buffalo	77	67	144	72.00
<b>2. Method of breeding</b>				
A. Natural	20	05	25	12.50
B. A.I.	36	37	73	36.50
C. Both	44	58	102	51.00
<b>3. Method of heat detection</b>				
A. Mucous discharge + bellowing	40	45	85	42.50
B. Mucous discharge + off feed	12	08	20	10.00
C. Mucous discharge + bellowing +mounting	18	22	40	20.00
D. Mucous discharge	30	25	55	27.50
<b>4. Expulsion of placenta</b>				
A. Within 1 hour	38	28	66	33.00
B. Within 2 hour	45	60	105	52.50
C. More than 2 hour	17	12	29	14.50
<b>5. Stage of heat which buffalo allowed for insemination/service</b>				
A. early heat	10	06	16	08.00
B. mid heat	78	86	164	82.00
C. later heat	12	08	20	10.00
<b>6. Pregnancy diagnosis</b>				
A. By experience farmers	31	25	56	28.00
B. By veterinary doctors	04	16	20	10.00
C. not done	65	59	124	62.00
<b>7. Age at first calving</b>				
A. Within 3 year	00	06	06	03
B. 3-4.5 year	49	66	115	57.5
C. 4.5-6 year	51	28	79	39.5
<b>8. Treatment of anoestrous/repeaters</b>				
A. Yes	77	86	163	81.5
B. no	23	14	37	18.5
<b>9. Breeding after calving</b>				
A. Within 3 months	11	36	47	23.50
B. 3-6 months	68	54	122	61.00
C. More than 6 months	21	10	31	15.50
<b>10. Calving interval</b>				
A. Less than 18 month	17	25	42	21.00
B. More than 18 month	83	75	178	79.00

All the respondents observed their buffaloes for heat symptoms regularly. It was found according to my village's study noted that buffaloes were allowed for insemination 82.00 per cent and 10.00 per cent of the respondents followed the practice in mid-heat and later heat, respectively, whereas only 08.00 per cent observed to detect heat in an early stage. This is the support of the study with Kumar *et al.* (2019a), Kumar *et al.* (2020)<sup>[8]</sup>, Singh (2018)<sup>[18]</sup>, Singh *et al.* (2019a), and Godara *et al.* (2018b)<sup>[5]</sup> found Mostly respondents inseminate at the later stage of heat.

It was noted that pregnancy diagnosis not done by most of the farmers 62.00 per cent in the villages and 28.00 per cent respondents were pregnancy diagnosis done by experienced farmers and only 10.00 per cent by the qualified veterinarian in the study area. The results are in line with the findings of Dar *et al.* (2017)<sup>[3]</sup>, Tanwar *et al.* (2012)<sup>[17]</sup>, Sreedhar *et al.* (2017a)<sup>[16]</sup>, Singh (2018)<sup>[18]</sup>, Malsawmdawngliana and Rahman (2016)<sup>[10]</sup>. It was contradicted the study found that most farmers' pregnancy diagnosis was done Khadda *et al.* (2017)<sup>[7]</sup>. It was observed from my study that most of the respondents were not observed pregnancy diagnosis by the respondents because of lack of knowledge and experience about the pregnancy respondents don't know how to detect pregnancy of the buffaloes.

The study found that most of the respondents 57.50 per cent animals takes 3-4.5 year time for first calving followed that more than 4.5-6 year time takes by the 39.50 per cent buffaloes and only 03.00 per cent buffaloes first calving within 3 years of age. These were in agreement with Dhaka *et al.* (2017)<sup>[4]</sup>, Meena *et al.* (2016)<sup>[11]</sup> first calve at 1288±122 day, Singh *et al.* (2016)<sup>[11]</sup>.

The result indicated that all the respondents treated their buffaloes for anoestrus, repeat breeding, and sometimes infertility problem. It was found that 81.50 per cent of respondents treated the anoestrus and 18.50 per cent of respondents not treated their animals in the villages. These results are in consonance with the findings of Khadda *et al.* (2017)<sup>[7]</sup>, Sivaji *et al.* (2018), and this observation highly contradicted by the Malsawmdawngliana and Rahman (2016)<sup>[10]</sup>.

It was observed that 61.00 per cent buffaloes successive breeding done 3-6 months after calving followed that 23.50 per cent breeding their animals within 3 months of calving and 15.50 per cent breeding done after 6 months of calving. Similar findings were reported by Kumar *et al.* (2020)<sup>[8]</sup>, Malsawmdawngliana and Rahman (2016)<sup>[10]</sup>, Singh (2018)<sup>[14]</sup>, Singh *et al.* (2019a), Yadav *et al.* (2016)<sup>[19]</sup>, Sekhar *et al.* (2017)<sup>[12]</sup>.

It was found in the study mostly 79.00 per cent of the respondent's buffaloes were taken more than 18 months of calving interval and only 21.00 per cent calving within 18 months of age. These were in agreement with Dhaka *et al.* (2017)<sup>[4]</sup>, Singh (2018)<sup>[18]</sup>, Sreedhar *et al.* (2017a)<sup>[16]</sup> found 73.75 percent were taking more than 18 months' time. Meena *et al.* (2016)<sup>[11]</sup> observed that the calving interval of buffaloes is 495±45. Calving interval is more in the buffaloes this is because of repeated breeding and due to lack of feeding management.

## Conclusion

It was noted that in the study most of the respondents of Kanpur Nagar prefer that both methods of insemination. Repeat breeding and anoestrus are highly affected to the dairy buffaloes this is because of hot weather in summer and poor

management practices used by the respondents. This is also observed that pregnancy diagnosis is not done by most of the respondents because of lack of awareness. It was found in the study mostly 79.00 per cent of the respondent's buffaloes were taken more than 18 months of calving interval because of lack of management practices adopted by the respondents in the district.

## Acknowledgement

With limitless humility, I bow my head to Almighty, Merciful Compassionate and Supreme power 'God' who showered his mercy on me and blessed me with the favorable circumstances to go through his gigantic task. I feel golden opportunity with great pleasure in acknowledging my profound sense of veneration and gratitude to my major advisor, Dr. Ved Prakash. The authors are thankful to the senior sir Dr. Raj Kumar, Department of Soil Conservation & Water Management, Chandra Shekhar Azad University of Agriculture & Technology, Kanpur, (U.P.) India for his keen interest, valuable guidance, and constructive criticism throughout the pursuit of the present research and vital suggestion during preparation of this manuscript.

## References

1. Anonyms. Annual report 2017-18. Department of Animal Husbandry, Dairying and Fisheries, GOI 2018, 2-3.
2. Chakravarthi Dr. M. Kalyana, Bharadwaja Dr. Marella, Kumar Kutty, Dr. Potu Ramesh Babu. Extent of Adoption of Improved Animal Husbandry Practices by Dairy Farmers in Kadapa District of Andhra Pradesh. International Journal of Science, Environment and Technology 2017;6(5):2815-2820.
3. Dar PA, Khan AA, Ahmad A, Qadri IA, Manzoor Ashiq, Khan Hilal Musadiq. Studies on Breeding and Feeding Practices of Buffalo in Vogue among Buffalo Farmers in Temperate Himalayan Region of Kashmir Valley. Indian Journal of Hill Farming, 2017, 83-87.
4. Dhaka BL, Meena GS, Meena NL. Reproductive Performance of Buffaloes under Field Conditions in Bundi District of Rajasthan, India. Int. J Curr. Microbiol. App. Sci 2017;6(4):595-599.
5. Godara Vikas, Gulati Harish Kr, Singh Narender Robin. Buffalo Breeding Management Practices Followed by Farmers of Western Haryana. J Krishi Vigyan 2018;6(2):13-16.
6. Iannuzzi L. Standard karyotype of the river buffalo (*Bubalus bubalis* L., 2n = 50). Cytogenetics and Cell Genetics 1994;67:102-113.
7. Khadda BS, Lata Kanak, Singh Brijesh, Kumar Raj. Study of buffalo husbandry practices in rural area of central Gujarat in India, Buffalo Bulletin 2017;36(1):75-87.
8. Kumar Avanish, Kumar Vipin, Upadhyay, Singh VP. Existing Dairy Husbandry Practices followed by Livestock owners in Farrukhabad District of Uttar Pradesh, India. Int. J Curr. Microbiol. App. Sci 2020;9(2):1863-1873.
9. Kumar J, Singh R, Dayal Somnath, Rajeshwar Singh, Hitesh and Singh Satyveer. Studies on feeding & breeding practices of dairy animal in western Uttar Pradesh. Journal of Pharmacognosy and Phytochemistry 2019, 29-36.
10. Malsawmdawngliana R, Rahman S. Management practices followed by the dairy farmers of Mizoram,

- India. *Journal of Livestock Science* 2016;7:220-225.
11. Meena BS, Verma HC, Singh Amit. Farmer's knowledge on Productive and Reproductive Performances of Buffalo under Small Holder Farming System. *Buffalo Bulletin* 2016;35:101-108.
  12. Sekhar MR, Rajanna N, Mahendra M, Chaudhary S. Comparative study of feeding and breeding management practices of dairy farmers in two different production systems. *Asian J. Dairy & Food Res* 2017;36(4):269-275.
  13. Singh R, Kumar P, Khandi SA, Bhadwal MS. Animal welfare practices followed by dairy farmers of Kathua district of Jammu and Kashmir State. *Journal of animal research* 2016;6(3):479-491.
  14. Singh R. Existing Animal Husbandry Practices in Narmada District of Gujarat in India. *International Journal of Animal Science and Technology* 2018;2(2):23-29.
  15. Sivaji DV, Natchimuthu K, Ramkumar S, Sreekumar D, Ganesan R. A Study on Adoption of Buffalo Husbandry Practices in Guntur and Prakasam Districts of Andhra Pradesh. *Res J Chem. Environ. Sci* 2018;6:35-39.
  16. Sreedhar S, Reddy Nagarjuna A, Sudhakar BV, Babu Ramesh P. Breeding management practices and Reproductive Disorders in Indigenous Cattles and Buffaloes. *Global Journal of bio-science and biotechnology* 2017;6(3):504-508.
  17. Tanwar PS, Kumar Yogendra, Sankhala Gopal. Study on Breeding and Feeding Management Practices Followed By Members and Non-members of Dairy Cooperatives in Jaipur District Of Rajasthan. *Indian J Dairy Sci* 2012;65(6):508-513.
  18. Viswkarma Singh Ruchi, Kushram Priyanka, Singh Sandeep Kumar and Sharma Shivangi. Existing status of buffalo husbandry practices in Jabalpur. *The Pharma Innovation Journal* 2018;7(2):08-11.
  19. Yadav SP, Paswan VK, Sawant P, Bhinchhar BK. Breeding and calf rearing management practices followed in Varanasi district of Uttar Pradesh, India. *Indian J Anim. Res* 2016;50(5):799-803.