



International Journal of Fauna and Biological Studies

Available online at www.faunajournal.com

I
J
F
B
S

International
Journal of
Fauna And
Biological
Studies

E-ISSN 2347-2677

P-ISSN 2394-0522

www.faunajournal.com

IJFBS 2021; 8(3): 38-42

Received: 22-03-2021

Accepted: 24-04-2021

Santa Kalita

(1) Department of
Environmental Science, Tezpur
University, Tezpur, Assam,
India

(2) Department of Zoology,
Gauhati University, Guwahati,
Assam, India

RK Bhola

Department of Zoology,
Gauhati University, Guwahati,
Assam, India

Efficacy of botanicals in controlling F₁ progeny production in *Tribolium castaneum* (Herbst)

Santa Kalita and RK Bhola

Abstract

Laboratory bioassays (30±2 °C temperature, 70±5% r. h. and 12:12, L: D conditions) were performed to evaluate the insecticidal effects of eight different plant extracts against *Tribolium castaneum* (Herbst). Dichloromethane extracts of *Lantana camara*, *Calotropis procera*, *Polygonum hydropiper*, *Jatropha curcas*, *Datura metel*, *Annona squamosa*, *Vitex negundo* and *Murraya koenigii* were applied to whole-wheat flour at different doses of 40, 50, 60 and 75 mg respectively and placed in different jars. Hundred newly emerged adults were introduced into each jar for mating, oviposition and later removed from the corresponding jars. The emerging adults (F₁) were counted and weighed. Results were analyzed statistically. It was observed that all the plant extracts significantly reduced the production of F₁ individuals (F₁ progeny production) with *Calotropis procera* and *Jatropha curcas* showing the highest efficacy.

Keywords: pest, botanicals, environment, control

1. Introduction

Tribolium castaneum (Herbst) (Coleoptera: Tenebrionidae) is a stored grain pest and is has a worldwide distribution. It is reddish brown in colour and is commonly known as the red flour beetle. It is a pest of cereals, flour, beans, pasta, pet food, dried flowers as well as museum specimens, chocolates, seeds and nuts (Via, 1999; Weston and Rattlingourd, 2000) [1, 2]. The egg to adult life cycle takes about 40-90 days and they may live to a period of three years (Rebecca and Thomas, 2003) [3]. Red flour beetles are prolific breeders. They continuously breed in warm areas round the year. The eggs, larvae, pupae, adults, their excreta and body parts contaminate the grains and make it unfit for consumption by man. They cause increase in moisture content and rise in temperature that leads to the production of molds and growth of toxigenic species (Magan *et al.*, 2003) [4]. Post- harvest losses due to insect infestations and secondary attack by bacteria and fungi in storage conditions account for 10% and 20% of the total production of cereals and legumes in developed and developing countries, respectively (Mason and McDonough, 2012; Nayak and Daglish, 2018) [5, 6]. Though chemical pesticides are used to control their population; these have many harmful effects on human health, environment and on non-target organisms. They have after-effects like pest resurgence and pesticide resistance that becomes a major problem in pest management. There are several environmental and safety concerns specially on use of chemicals like methyl bromide and dichlorvos, which have been in use for insect pest eradication. Hence, there is a prime need for evaluation and development of new insecticidal formulations for the control of stored product insect pests (Daglish *et al.*, 2018) [7].

Plants provide prospective alternatives to currently used toxic chemical insecticides. Plants contain bioactive compounds; these constitute a defence system in plants to protect them against attack of insects and other diseases. The bioactivities associated with the secondary metabolites present in plants counteract the activities of varied species of insects (Isman, 2006, 2008) [8, 9]. Thus, they can suffice as control measure against insect pest populations. Extracts derived from plants are locally and easily available, less expensive, biodegradable, have diverse modes of action, and have low toxicity to non-target organisms. They can be incorporated into integrated pest management systems and help in contributing to sustainable agricultural production (Geraldin *et al.*, 2020) [10].

Toxicity was observed in insect pests of storage like *T. castaneum* (Herbst), *Orzyaepphilus surinamensis* (L.), *Sitophilus oryzae* as well as *Rhyzopertha dominica* upon application of the essential oil of spices like anise (*Pimpinella anisum* L.) and peppermint (*Mentha piperita* L.)

Corresponding Author:

Santa Kalita

(1) Department of
Environmental Science, Tezpur
University, Tezpur, Assam,
India

(2) Department of Zoology,
Gauhati University, Guwahati,
Assam, India

(Shaaya *et al.*, 1991) ^[11]. Also, the essential oil of garlic (*Allium sativum*) showed insecticidal properties upon application on *T. castaneum* and *Sitophilus zeamais* (Motschulsky) (Ho *et al.*, 1996) ^[12]. Similarly, essential oil extracted from seeds of nutmeg (*Myristica fragrans* Houtt) (Huang *et al.*, 1997) ^[13] and bark of cinnamon (*Cinnamomum aromaticum* Nees) (Huang and Ho, 1998) ^[14] exhibited toxicity against *S. zeamais* and *T. castaneum*. Extracts of *L. camara* reduced the emergence of adults, induced delay in development and produced lower fecundity in *R. dominica* (Singh *et al.*, 1996) ^[15]. Leaf extracts of *J. curcas*, *L. camara*, *C. procera* and *D. metel* exhibited toxicity against *T. castaneum* (Kalita and Bholia, 2014) ^[16]. Twenty-three different plant extracts were evaluated for their efficacy on *Callosobruchus maculatus* (Boeke *et al.*, 2004) ^[17]. Promising results were found when pea flour extract was enriched with protein and used against *S. oryzae* to protect rice in storage (Kumar *et al.*, 2004) ^[18].

The present study was taken up to observe how the eight plant extracts affected the F₁ progeny production in the tested insect *T. castaneum*.

2. Materials and Methods

2.1 Insect rearing and preparation of plant extracts

T. castaneum was cultured in the laboratory at ambient conditions of temperature, relative humidity and photoperiod of (30±2°C), (70±5%) and (12:12, L:D conditions) respectively. Leaves of *Lantana camara*, *Calotropis procera*, *Polygonum hydropiper*, *Jatropha curcas*, *Datura metel*, *Annona squamosa*, *Vitex negundo* and *Murraya koenigii* were collected, washed and air-dried for three to five days, followed by drying in the oven at 40°C for 24 hours after which they were powdered in an electric blender. The extraction was carried out in Soxhlet apparatus using dichloromethane. The extract was collected, air-dried to vaporize the solvent and covered by aluminum foil. This was kept at 4°C and stored for conduction of the bioassays.

2.2 Application of the extracts

100g whole-wheat flour was sterilized in an oven for 12h at 30°C. Thereafter, 40, 50, 60 and 75 mg of the extracts of the tested plants were added to the sterilized flour for each bioassay respectively. The flour was then shifted to jars measuring 500 ml. In each jar, hundred newly emerged adults were placed; then kept in an incubator for mating and oviposition at a temperature of 30°C for 4 days. The flour was then sieved to remove the adults and placed in different jars as per the treatments applied. The mouths of the jars were tied with muslin cloth and kept at a temperature of 30°C for emergence of adults. The F₁ adults were counted and weighed after complete emergence. Five replicates were taken for each treatment and one control was set up. Results were statistically analyzed.

3. Results

Extracts from eight plants were applied to the parent insect *Tribolium castaneum* in order to evaluate their effect(s) on F₁ generation. It was observed that in all the concentrations, the tested plants exhibited significant reduction in the number of adults of F₁ generation. The bioassay using dichloromethane extract of the eight plants on production of F₁ generation is presented in Table 1. *C. procera* (42.7-58.8) exhibited highest effectiveness with the production of least number of progeny followed by *J. curcas* (41.4-66.6), *P. hydropiper* (54.3-56.1),

D. metel (43.2-69.4), *V. negundo* (55.5-62.4), *L. camara* (55.2-68.4), *A. squamosa* (57.9-65.2) and *M. koenigii* (61.9-70). As the concentration increased from 40mg to 75mg, there was decrease in the number of F₁ adults' emergence. The highest number of F₁ progeny was in the control bioassay with a production of about 95.5 individuals, which was much higher than any of the treated bioassays.

The weight of the F₁ progeny with regard to a single insect did not vary significantly as compared to the control in all the bioassays.

3.1. Effect of the extracts at a concentration of 40 mg on F₁ progeny

For all the plant extracts tested, the numbers of F₁ progeny produced were less than that of the control. *P. hydropiper* showed the highest efficacy with an average emergence of 56.1 individuals against control, which showed an emergence of 95.4 number of individuals of *T. castaneum*. The order of efficacy was *P. hydropiper* > *C. procera* > *V. negundo* > *A. squamosa* > *J. curcas* > *L. camara* > *D. metel* > *M. koenigii* with an emergence of 56.1, 58.8, 62.4, 65.2, 66.6, 68.4, 69.4 and 70 numbers of *T. castaneum* individuals respectively. Thus, all the tested extracts could reduce the production of F₁ individuals significantly. However, there was no significant reduction in weight of the emerged individuals of F₁ progeny and was almost same as that of control.

3.2. Effect of the extracts at a concentration of 50 mg on F₁ progeny

C. procera at a concentration of 50mg showed the highest efficacy in controlling F₁ progeny production. It led to the emergence of 54.4 number of individuals as compared to 95.4 in control. The order of efficacy was *P. hydropiper* > *J. curcas* > *V. negundo* > *D. metel* > *A. squamosa* > *L. camara* > *M. koenigii* with 55.8, 58.4, 59.9, 61.6, 63.2, 65.4, 68.7 average numbers of emerging individuals respectively. There was no significant reduction in weight of the emerged individuals of the F₁ progeny. But in the treatment with *L. camara*, the weight was found to be 1.7mg against 2.25 in control; however, no significant differences in weight could be observed.

3.3. Effect of the extracts at a concentration of 60 mg on F₁ progeny

J. curcas showed the highest efficacy in controlling the emergence of F₁ individuals at 60mg concentration. Average weight of emerged individual was 1.58mg against 2.25mg in control. However, it was not significantly different from control. The order of efficacy was *J. curcas* > *C. procera* > *D. metel* > *P. hydropiper* > *V. negundo* > *L. camara* > *A. squamosa* > *M. koenigii* with emergence of an average of 42.2, 45.0, 46.2, 54.0, 57.0, 57.4, 61.8, 65.8 number of individuals in F₁ progeny against 95.4 numbers in control.

3.4. Effect of the extracts at a concentration of 75 mg on F₁ progeny

The lowest emergence of F₁ individuals was in the bioassay treated with 75mg extract of *J. curcas*. The order of efficacy was observed to be *J. curcas* > *C. procera* > *D. metel* > *L. camara* > *P. hydropiper* > *V. negundo* > *A. squamosa* > *M. koenigii* with emergence numbers of an average of 41.4, 42.7, 43.2, 54.3, 55.2, 55.5, 57.9, 61.9 individuals which is significantly different from control where the observed number were 95.5 individuals. Though there was no

significant difference in weight, however, the bioassays treated with *J. curcas* and *L. camara* resulted in production of individuals weighing 1.56mg and 1.66mg respectively against

the control, which was 2.25mg.

Tables and Figures

Table 1: Effect of dichloromethane (DCM) plant extract on F₁ progeny

Plant	Concentration							
	40 mg/100g wheat Flour		50 mg//100g wheat Flour		60 mg//100g wheat Flour		75 mg/100g wheat Flour	
	No. of F1 adults	Wt. of F1 progeny (mg)	No. of F1 adults	Wt. of F1 progeny (mg)	No. of F1 adults	Wt. of F1 progeny (mg)	No. of F1 adults	Wt. of F1 progeny (mg)
Control	95.4±3.65	215.6 (2.25)	95.4±3.6	215.6 (2.25)	95.4±3.6	215.6 (2.25)	95.4±3.6	215.6 (2.25)
<i>C. procera</i>	58.8±5.4*	130.4±5.2 (2.21)	54.4±2.8*	106.6±7.1 (1.96)	45.0±5.2*	83.2±9.9 (1.80)	42.7±5.4*	79.3±8.3 (1.86)
<i>J. curcas</i>	66.6±4.22*	138.2±10.4(2.07)	58.4±7.1*	111.4±14.1(1.91)	42.2±7.1*	66.8±11.2(1.58)	41.4±4.8*	64.4±13.1(1.56)
<i>P. hydropiper</i>	56.1±5.1*	128.4±8.9(2.29)	55.8±5.3*	126.80±9.9(2.27)	54.0±5.4*	115.1±6.9(2.14)	54.3±7.0*	118.7±7.3(2.18)
<i>D. metel</i>	69.4±4.39*	124.2±7.7(1.78)	61.6±5.1*	123.2±16.3(2.0)	46.2±6.3*	86.8±12.5(1.88)	43.2±3.7*	83.8±9.4(1.93)
<i>V. negundo</i>	62.4±4.1*	126.0±10.1(2.1)	59.9±3.2*	124.4±12.1(2.08)	57.0±4.8*	119.4±11.8(2.09)	55.5±4.9*	114.0±6.2(2.05)
<i>L. camara</i>	68.4±5.94*	123.2±14.2(1.80)	65.4±5.4*	111.0±9.3(1.7)	57.4±9.9*	93.0±16.5(1.62)	55.2±6.1*	91.5±12.5(1.66)
<i>A. squamosa</i>	65.2±3.7*	136.6±12.3(2.09)	63.2±4.8*	134.0±12.3(2.12)	61.8±6.1*	130.0±15.4(2.10)	57.9±5.6*	125.7±5.7(2.17)
<i>M. koenigii</i>	70.0±6.20	140.2±7.8(2.0)	68.7±6.7*	135.2±8.2(1.97)	65.8±3.5*	124.6±7.9(1.89)	61.9±3.6*	118.1±9.1(1.90)

(Figure in parenthesis represent the average weight of *T. castaneum*.)

Average weight of single insect (of F₁ progeny) at varied concentrations of same/ different plant extract do not differ significantly

*Significantly lower than control)

4. Discussion

The results showed that the *T. castaneum* population level were lower in the treated wheat samples as compared to the ones in the control. Similar results were observed on several insect pests using varied plant extracts. Extracts of *Peganum harmala*, *Ajuga iva* and *Aristolochia baetica* inhibited F₁ progeny production in *T. castaneum* (Jbilou *et al.*, 2008) [19]. *P. harmala* seeds contain bioactive compounds that cause developmental, physiological and histopathological effects in *T. castaneum*. Methanol extracts of *Peganum harmala* (L.) seeds when mixed with food led to inhibition of larval growth, reduced food intake, increased duration of larval period and inhibition of F₁ progeny production in *T. castaneum* (Jbilou and Sayah, 2008) [20]. Extracts from plants and their oils contain camphor, eugenol, 1, 8-cineole which have properties that can result in inhibiting production of progeny and causing mortality in insect pests (Obeng-Ofori and Reichmuth, 1997) [21]. Inhibition of adult emergence was also reported in *Sitophilus granaries* and *Rhyzopertha dominica* treated with gounded seeds of black pepper (*Piper nigrum* L.) and fruit powder of red pepper (*Capsicum annum* L.) (Shabnam and Nouraddin, 2010; Singh *et al.*, 2009) [22, 23]. *Calotropis procera* significantly reduced the number of adult emergence of *R. dominica* Fabr. (Jacob and Sheila, 1993) [24]. All the eight plant extracts exhibited significant reduction in the number of F₁ adults of *T. castaneum* at the different tested concentrations. *C. procera* was the most effective, which resulted in the production of least number of F₁ progeny. The order of efficacy was *C. procera* > *J. curcas* > *P. hydropiper* > *D. metel* > *V. negundo* > *L. camara* > *A. squamosa* > *M. koenigii*. The adult emergence of *C. procera* was in the range of 58.8, 54.4, 45 and 42.7 against 40, 50, 60 and 75 mg/ 100 gram flour respectively. As the concentration increased from 40mg to 75mg, the emergence of F₁ individuals reduced in number. A similar study revealed the potency of *Piper nigrum* powder on *Callosobruchus maculatus*. It reduced oviposition and adult emergence in a significant manner (Rajapakes, 1990) [25]. The two varieties of pepper (black pepper and red pepper) helped to reduce the emergence of adults of *Callosobruchus chinensis* L. (Morrallo – Rejesus *et al.*, 1990) [26]. Both larvae and adult development of *T. castaneum* in

grains of millet was reduced by using clove and West African black pepper (Lale and Ajayi, 2000) [27]. Edible fatty oils of *Moringa oleifera*, *Sesamum indicum* and *Olea europaea* stopped progeny emergence of *T. castaneum* in broken millet (Appert, 1987) [28]. F₁ progeny production in *T. castaneum* was totally inhibited by nutmeg oil (Huang *et al.*, 1997) [13] and by *Elettaria cardamomum* (L.) oil (Huang *et al.*, 2000) [29] when applied to the diet. Azadirachtin significantly reduced offspring production in *T. confusum* (Athanasios *et al.*, 2005) [30]. There was observation of F₁ inhibition in *Zabrotes subfasciatus* upon application of *Chenopodium ambrosioides*, *A. indica* and *Datura stramonium* in powder form (Araya and Eman, 2009) [31].

The present study showed that there was no significant difference in the weight of single insect of the F₁ generation at diverse concentrations of the extracts of the same as well as the different plants tested. However, in the bioassays using *J. curcas* and *L. camara* extracts, there was observance of a slight decrease in the weight of emerging individuals. Nicotine application in *Heliothis virescens* produced lesser number of survivors, decrease in weight of pupae and longer developmental period (Gunasena *et al.*, 1990) [32]. F₁ progeny production was inhibited in *C. maculatus*, by applying *Gnidia kaussiana* Meisn extracts (Kosini and Nukene, 2017) [33]. *Hemizygia welwitschii* extracts were toxic against adults of *C. maculatus* and significantly inhibited emergence of the F₁ progeny in treated grains (Tagne *et al.*, 2019) [34].

5. Conclusion

The extracts of the tested plants exhibited positive results in reducing the number of F₁ individuals significantly. *C. procera* and *J. curcas* showed highest efficacy producing lesser number of F₁ progeny followed by *P. hydropiper*. The extracts of *D. metel* was more effective than *V. negundo*, while *L. camara* showed more potency than *A. squamosa* and *M. koenigii*. However, all the plant extracts showed good efficacy as compared to the control thus revealing the possibility of their use for controlling the population of *T. castaneum*. The isolation and identification of the active ingredients from botanicals and the mode of action on insect pests can be used for combating pests of food commodities,

specifically in storage. These botanicals are eco-friendly, locally available and cheaper than their chemical counterparts. They pose low risk hazards and have less residual toxicity on non-target organisms. Their isolation and use to combat insect pests can usher in an era of health and environment friendly insecticides.

6. References

- Via S. Cannibalism facilitates the use of a novel environment in the flour beetle, *Tribolium castaneum*. *Heredity* 1999;82:267-275. <https://doi.org/10.1038/sj.hdy.6884820>
- Weston PA, Rattlingourd PA. Progeny production by *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Oryzaephilus surinamensis* (Coleoptera: Silvanidae) on maize previously infested by *Sitotroga cerealella* (Lepidoptera: Gelechiidae). *Journal of Economic Entomology* 2000;93(2):533-535. <https://doi.org/10.1603/0022-0493-93.2.533>
- Rebecca B, Thomas RF. "Confused Flour Beetle, *Tribolium confusum* and Red Flour Beetle, *Tribolium castaneum*". Entomology and Nematology Department, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, 2003.
- Magan N, Hope R, Cairns V, Aldred D. Postharvest fungal ecology: impact of fungal growth and mycotoxin accumulation in stored grain. *Eur. J Plant Pathol* 2003;109:723-730. DOI: 10.1023/A:1026082425177
- Mason LJ, McDonough M. Biology, behavior, and ecology of stored grain and legume insects. In: Hagstrum D. W, Phillips T. W, Cuperus G. (Ed.), *Stored Product Protection*. Kansas State University, Manhattan, KS 2012, 7-20.
- Nayak MK, Daglish GJ. Importance of Stored Product Insects. In: Athanassiou C., Arthur F. (Ed.), *Recent Advances in Stored Product Protection*. Springer, Berlin, Heidelberg, 2018, 1-17. https://doi.org/10.1007/978-3-662-56125-6_1
- Daglish GJ, Nayak MK, Arthur FH, Athanassiou CG. Insect Pest Management in Stored Grain. In: Athanassiou C., Arthur F. (Ed.), *Recent Advances in Stored Product Protection*. Springer, Berlin, Heidelberg, 2018, 45-63. https://doi.org/10.1007/978-3-662-56125-6_3
- Isman MB. Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Ann. Rev. Entomol* 2006;51:45-66. DOI: 10.1146/annurev.ento.51.110104.151146
- Isman M B. Botanical insecticides: for richer, for poorer. *Pest Manage. Sci* 2007;64:8-11. <https://doi.org/10.1002/ps.1470>
- Geraldin MW, Lengai James WM, Ernest Mbega R. Phytochemical activity and role of botanical pesticides in pest management for sustainable agricultural crop production, *Scientific African* 2020;7:1-13. <https://doi.org/10.1016/j.sciaf.2019.e00239>.
- Shaaya E, Ravid U, Paster N, Juven B, Zisman U, Pissarev V. Fumigant toxicity of essential oils against four major stored product insects. *J Chem. Ecol* 1991;17:499-504.
- Ho S H, Koh L, Ma Y, Huang Y, Sim KY. The oil of garlic, *Allium sativum* L. (Amaryllidaceae), as a potential grain protectant against *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. *Postharvest Biol. Technol* 1996;9:41-48.
- Huang Y, Tan JM WL, Kini RM, Ho SH. Toxic and antifeedant action of nutmeg oil against *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. *J. Stored Prod. Res* 1997;33(4):289-298. [https://doi.org/10.1016/S0022-474X\(97\)00009-X](https://doi.org/10.1016/S0022-474X(97)00009-X)
- Huang Y, Ho SH. Toxicity and antifeedant activities of cinnamaldehyde against the grain storage insects, *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. *Journal of Stored Products Research* 1998;34(1):11-17. DOI: 10.1016/S0022-474X(97)00038-6.
- Singh H, Mrig KK, Mahla JC. Effect of different plant products on the fecundity and emergence of lesser grain borer *Rhyzopertha dominica* wheat grains. *Annals of Biology* 1996;12(1):96-98.
- Kalita S, Bhola RK. Repellency and toxicity of some plant extracts against *Tribolium castaneum* (Herbst). *Global Journal for Research Analysis* 2014;3(6):8-9.
- Boeke SJ, Barnaud C, van Loon JA, Kossou DK, van Huis A, Dicke M. Efficacy of plant extracts against the cowpea beetle, *Callosobruchus maculatus*. *Int. J. Pest Man* 2004;50:251-258. <https://doi.org/10.1080/09670870410001702206>
- Kumar S. Occupational Exposure and reproductive dysfunction. *J Occup Health* 2004;46:1-19. <https://doi.org/10.1539/joh.46.1>
- Jbilou R, Ennabili A, Sayah F. Insecticidal activity of four medicinal plant extracts against *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *African Journal of Biotechnology* 2006;5:936-940.
- Jbilou R, Sayah F. Effects of *Peganum harmala* (Zygophyllaceae) seed extracts on the development of *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Int J Trop Insect Sci* 2008;27(3-4):199-209. DOI: 10.1017/S1742758407850971
- Obeng-Ofori D, Reichmuth CH. Bioactivity of eugenol, a major component of essential oil of *Ocimum suave* (Wild.) against four species of stored-product Coleoptera. *Int. J Pest Manage* 1997;43:89-94. <https://doi.org/10.1080/096708797229040>
- Shabnam A, Nouraddin S. Insecticidal activities of two powdered spices, black pepper and red pepper on adults of *Rhyzopertha dominica* (F.) and *Sitophilus granaries* (L.). *Mun. Entomology & Zoology* 2010;5(2):600-607.
- Singh N, Sharda MP, Francis OS. Evaluation of important triterpenes in field and *in vitro* plants of *Syzygium cordatum*, *J Environ. Res. Develop* 2009;3(3):601-606.
- Jacob S, Sheila MK. A note on the protection of the stored rice from the lesser grain borer, *Rhyzopertha dominica* Fabr. by indigenous plant products. *Indian J. Entomol* 1993;55(3):337-339.
- Rajapakes RHS. Effects of five botanicals as protectants of green gram against the pulse beetle, *Callosobruchus maculatus* In: Fujii (Ed.), *Bruchids and Legumes: Economics, Ecology and Co-evolution*. Kulwer Academic Publisher, Netherland, 1990, 85-90.
- Morallo-Rejesus B, Maini HA, Ohasawa K, Yamato I. Insecticidal actions of several plants to *Callosobruchus chinensis* L. In: Fujii, (Ed.), *Bruchids and Legumes: Economics, Ecology and Co-evolution*. Kulwer Academic Publisher, Netherland, 1990, 97-100.
- Lale NES, Ajayi F. Suppressing infestation of stored pearl millet *Pennisetum glaucum* (L) R. Br. by *Tribolium*

- castaneum* (Herbst.) with insecticidal essential oils in Maiduguri, Nigeria. *Journal of Plant Diseases and Protection* 2000;107(4):392-398.
28. Appert J. *The Storage of Food: Grains and Seeds*. London, Macmillan Publishers, 1987, 146.
 29. Huang Y, Lam SL, Ho SH. Bioactivities of essential oil from *Elletaria cardamomum* (L.) Maton. to *Sitophilus zeamais* Motschulsky and *Tribolium castaneum* (Herbst), *Journal of Stored Products Research* 2000;36(2):107-117. DOI: 10.1016/S0022-474X(99)00040-5
 30. Athanassiou CG, Kontodimas DC, Kavallieratos NG, Veroniki MA. Insecticidal effect of NeemAzal against three stored product beetle species on rye and oats. *J. Econ. Entom* 2005;98(5):1733-1738. <https://doi.org/10.1603/0022-0493-98.5.1733>
 31. Araya G, Eman D. Evaluation of botanical plants powders against *Zabrotes subfasciatus* (Boheman) (Coleoptera: Bruchidae) in stored haricot beans under laboratory condition. *African Journal of Agricultural Research* 2009;4(10):1073-1079. <https://doi.org/10.5897/AJAR.9000308>
 32. Gunasena GH, Vinson SB, Williams HJ. Effects of Nicotine on growth, development and survival of the tobacco budworm (Lepidoptera :Noctidae) and the parasitoid, *Campoletis sonorensis* (Hymenoptera : Ichneumonidae). *J Econ. Ent* 1990;83(5):1777-1782. <https://doi.org/10.1093/jee/83.5.1777>
 33. Kosini D, Nukenine EN. Bioactivity of novel botanical insecticide from *Gnidia kaussiana* (Thymeleaceae) against *Callosobruchus maculatus* (Coleoptera: Chrysomelidae) in stored *Vigna subterranea* (Fabaceae) grains. *J Insect Sci* 2017;17(1):311-7. doi: 10.1093/jisesa/iex004
 34. Tagne GF, Haman KT, Jean PA, Narcis T, Calvin MZ, Cornel A. Control of *Callosobruchus maculatus* (Coleoptera: Chrysomelidae) using fractionated extracts from Cameroonian *Hemizygia welwitschii* (Lamiaceae) leaf on stored *Vigna unguiculata* (Fabales: Fabaceae). *Journal of Insect Science* 2019;19(2):22. <https://doi.org/10.1093/jisesa/iez029>