



E-ISSN 2347-2677

P-ISSN 2394-0522

[www.faujournal.com](http://www.faujournal.com)

IJFBS 2021; 8(3): 11-17

Received: 03-03-2021

Accepted: 06-04-2021

**Anju Bharti**

Biopesticides and Toxicological  
Lab., Department of Zoology,  
D.B.S. College, Kanpur, Uttar  
Pradesh, India

**Indrani Dubey**

Biopesticides and Toxicological  
Lab., Head Department of  
Zoology, D.B.S. College, Kanpur,  
Uttar Pradesh, India

**BS Chandel**

Biopesticides and Toxicological  
Lab., Department of Zoology,  
D.B.S. College, Kanpur, Uttar  
Pradesh, India

**Corresponding Author:**

**Anju Bharti**

Plant Protection Laboratory,  
Department of Zoology, D.B.S.  
College, Kanpur, Uttar Pradesh,  
India

## Phago-repellent biopotency of *Cichorium intybus*, *Inula racemosa*, *Tagetes minuta* and *Chrysanthemum cinerariaefolium* as herbal alternatives to synthetic insecticides against *Earias vittella* Fabricius (Lepidoptera: Noctuidae) on Okra in Kanpur region

Anju Bharti, Indrani Dubey and BS Chandel

DOI: <https://doi.org/10.22271/23940522.2021.v8.i3a.824>

### Abstract

To evaluate the effectiveness of selected asteraceous botanical extractives in management of spotted bollworm, *Earias vittella* Fabricius (Lepidoptera: Noctuidae) an experiment was conducted in the plant product laboratory, Department of Zoology, D.B.S. College, Kanpur Nagar during 2021. Phago-repellent effectiveness of ten indigenous naturally occurring asteraceous plant extracts viz., aerial parts of *Cichorium intybus* (L.), *Chromolaena odorata* Linn., *Chrysanthemum cinerariaefolium* (trev.) Vis., *Inula racemosa* Hook. F., *Mantisalca duriaeri* Birq. Et Cavill., *Reichardia tingitana* (L.) Roth, *Rhaponiticum acaule* (L.) DC, *Scorzonera undulate* Vahl, *Spilanthes paniculata* Well ex DC and *Tagetes minuta* Linn. were prepared under the laboratory conditions. For testing the phago-repellent effect, the okra leaves were used as food against larvae of *Earias vittella*. In each set of extract and one control was introduced, where the leaves pieces were dipped in Benzene + emulsified water only. The okra leaf pieces fastened under clip and left under electric fan for about thirty minutes, so as to complete dry up the extract. The treated foods were kept in jar (23cm x 10cm) on moist filter paper. Then third instars, 24 hours starved larvae of *E. vittella* were released in each jar. After four hours of the release of larvae the data was collected on the number of larvae reached and repelled at each treated food. Three replication of treatment were made. The data depicted from the results in the present experiment of repellent test against larvae of *Earias vittella*. All the plant extracts showed a good repellent property, when compared with the control. The repellent property was observed based on their EC<sub>50</sub> values as: *C. intybus* (0.0823) > *I. racemosa* (0.1043) > *T. minuta* (0.1620) > *C. cinerariaefolium* (0.1677) > *M. duriaeri* (0.1788) > *S. undulate* (0.3127) > *A. paniculata*, (0.3805) > *R. acaule* (0.4405) > *R. tingitana* (0.6368) > *C. odorata* (0.8870), respectively. The relative EC<sub>50</sub> values of the above extracts are in descending order, 10.7700 > 8.5043 > 5.4753 > 5.0227 > 4.9608 > 2.8365 > 2.3311 > 1.3929 > 1.0000 times as repellents, where as *C. odorata* is taken as a unit.

**Keywords:** *Earias vittella*, *Cichorium intybus*, *Inula racemosa*, *T. minuta*, Bhindi (Okra)

### 1. Introduction

Bhindi, *Abelmoschus esculentus* (Linn.) Moench commonly known as okra is an important summer vegetable crop grown all over Indian sub Zoogeographical region. Okra is nutritious vegetable which plays an important role to meet the demand of vegetable of the country when vegetable are scanty in the market. Okra contains large quantities of carbohydrate, potassium, vitamin B, vitamin C, protein, folic acid and calcium. It's low in calories and has high dietary fiber content. Okra is a nutritious vegetable which plays an important role to meet the demand of vegetables of the India when vegetable are scanty in the market. Okra mucilage is suitable for medicinal and industrial application (Radake and Undirwadem 1981, Kumar *et al.* 2013) <sup>[1, 2]</sup>.

The spotted bollworm, *Earias vittella* Fabricius (Lepidoptera: Noctuidae) is a serious pest of vegetables ie. Okra, eggplant, pepper melon, cucumber, pumpkin and cotton. The crop is vulnerable to attack of many insect pests, among them *Earias vittella* Fabricius are the most important pest causing direct damage to the marketable fruits (Konar and Rai, 1990) <sup>[3]</sup>. *Aphis gossypii* and *Earias vittella* is alone reported to cause 57.1% fruit infestation and 54.04% yield loss in okra (Chaudhary and Dadheech, 1989) <sup>[4]</sup> Leaves can be damaged to such an extent that they wilt and fall off.

This impairs photosynthesis, thereby weakening the plant even further, and may render fruits unmarketable.

The synthetic chemical insecticides are very effective and used in okra insect pest management for a long time and to ensure higher crop and vegetable yield. The excessive and injudicious use of these chemical pesticides led to many problems like development of resistance, induction of resurgence, environment pollutions and human health (Park *et al.* 2011, Chandel *et al.* 2018) [5, 6]. The side effects have forced to look for naturally occurring ecofriendly indigenous asteraceous herbal alternatives to chemical insecticides especially for vegetables like okra where fruits are plucked at an interval of every 2-3 days.

Various plant extractives have been studied for insecticidal activity globally (Dubey *et al.* 2004, Chandel and Sing) [7, 8] and majority of them are insect feeding deterrent (Dang *et al.* 2010, Antonius and Hagazy 1987) [9, 10]. More than 140 compounds, which are chemically diverse and structurally complex, have been isolated from the leaves, seed oil and bark of neem (Ahmed *et al.* 1999 and Al-Lavati *et al.* 2002) [11, 12]. Phytochemicals are often distasteful and toxic to many agricultural insect-pests. They can modify feeding deterrence behaviour of an insect (Schmidt and Strelake, 1994) [13]. Moreover, as most spotted bollworm species have become resistant to many insecticidal agents (Pavela, 2006, Rashid *et al.* 2013) [14, 15], managing these pests in greenhouses and in the field is becoming problematic (Srivastava and Awasthi 1958, Mendes, *et al.* 1997, Kim *et al.* 2011) [16, 17, 18]. Therefore, in the present investigation the selected asteraceous naturally occurring plant extractives are efficient and environmentally friendly pest control alternatives must be developed to replace synthetic insecticides.

## 2. Materials and Methods

Experiments were conducted in the Entomology Research plant product Laboratory and experimental field of Department of Zoology, D.B.S. College, Kanpur, U.P., India. Geographically, the districts Kanpur is located in between latitudes 25.26° and 26.58° North and longitudes 19.31° and 84.34° East, Kanpur is situated at an elevation of about 127.117° metres above the mean sea level and has a semi-arid subtropical climatic conditions.

### 2.1 Procurement of raw herbal materials

The plants parts used for extracts were surveyed, identified and collected mainly from wasteland and wild areas and some plants were collected from cultivated fields of the farmers. The investigations on the screening of various available indigenous naturally occurring ten asteraceous plant extracts viz., aerial parts of *Cichorium intybus* (L.), *Chromolaena odorata* Linn., *Chrysanthemum cinerariaefolium* (trev.) Vis. *Inula racemosa* Hook. F., *Mantisalca duriaeri* Birq. Et Cavill., *Rechardia tingitana* (L.) Roth, *Rhaponticum acaule* (L.) DC, *Scorzonera undulate* Vahl, *Spilanthes paniculata* Well ex DC and *Tagetes minuta* Linn. Were screened for their repellent biopotency against spotted bollworm, *Earias vittella* under laboratory trials.

### 2.2 Process of Herbal Powder Extraction

Fresh collected ten plant parts (leaves, Flowers and seeds, rhizomes etc) were washed with distilled water and kept in the laboratory for 7 days for air drying followed by one day sun

drying before making powder. Electric grinder was used to have coarse powder then these were passed through a 60-mesh sieve to get fine powder. Powders were kept in polythene bags at room temperature and properly sealed to prevent quality loss. For the extraction, Soxhlet Apparatus was used; about 20g powder of each category of powder were extracted with 300 ml of different solvents (n-hexane, acetone, methanol, petroleum ether and distilled water). Extraction of each category of powder was done in about 12 hrs. After soxhlet extraction, the material was run on rotary evaporator (Jilani, *et al.* 1985, Abubakar *et al.* 2000, Ramamurthy, *et al.* 2002) [19, 20, 21]. The extracts were concentrated on rotary evaporator by removing the excess solvent under vacuum. After evaporation of solvent with rotary evaporator the remaining extracted material was kept on water bath for removing remaining solvent from the extracts. The extracts were stored at 4°C prior to application.

### 2.3 Stock Solution Preparation and Formulations

For stock solution, 50ml. extract in each case was taken into reagent bottles and 50ml. benzene was added in it to dissolve the constituents of the materials. The mouth of the bottles were stopper with airtight corks after which, these bottles containing the solutions were kept in refrigerator. Three concentrations (0.25, 0.5, 1.0, 1.5, 2.0 percent) were used for experiments on repellent tests in the laboratory conditions. However, only three concentrations (0.5, 1.0 and 2.0 percent) were used for insecticidal test in the laboratory. The different concentrations of the herbal extracts were prepared from the stock solution using benzene as solvent and Triton X-100 as emulsifier. The level of solvent and emulsifier were kept constant (Kamaraj *et al.* 2011, Chandel and Sengar, 2018) [22, 23].

### 2.4 Experimental Equipments

Small plastic jars (capacity 50 ml) were used for the experiment, there was one set of two jars joined by clear plastic pipe of 1.0 cm diameter at an angle of 180 degree for each replication. One jar of each set was provided with 10 g of grains given the name 'A' while the other jar was kept empty and given the name 'B'. In jar 'A', the grains treated with extracts were placed, while the jar B remained empty. The jars used for experiment were disinfected with alcohol.

## 3. Experimental Process

The Phago-repellent test was carried out under laboratory condition against nymph and adults of *A. gossypii*. The okra leaves were used as food for the spotted bollworm, *Earias vittella* larvae. Leaves were treated with different concentrations (0.25, 0.5, 1.0, 1.5 and 2.0 per cent) for two minutes. The treated leaves were left under electric fan for about half an hour, to make a dry film of the extracts on the leaves for each set of extract and one control. The treated foods were kept in jar (23cm x 10cm) on moist filter paper. The untreated leaves were dipped in Benzene + emulsified water only. Ten starved *E. vittella* larvae were released in each jar along with control. Three replicates per treatments were maintained. The treated jars either repelled the insects or forced them to move from treated jars (T) to an empty jar (C) through the plastic pipe. The ones found in plastic pipe were considered repelled individuals. Number of *E. vittella* larvae reached to the food or repelled to untreated empty jar, were recorded.

**Table 1:** List of selected asteraceous plants materials used for extraction

S. No.	Scientific Name	Common names	Part Used	Family
1.	<i>Acmella paniculata</i> Well ex DC	Toothache Plant	Leaves	Asteraceae
2.	<i>Chromolaena odorata</i> Linn.	Siam weed	Leaves	Asteraceae
3.	<i>Chrysanthemum cinerariaefolium</i> (trev.) Vis.	Insecticide Daisy	Roots	Asteraceae
4.	<i>Cichorium intybus</i> (L.)	Chicory	Aerial parts	Asteraceae
5.	<i>Inula racemosa</i> Hook. f	Puskarmul	Roots	Asteraceae
6.	<i>Mantisalca duriaeri</i> Birq. Et Cavill.	Spach	Flowers	Asteraceae
7.	<i>Rechardia tingitana</i> (L.) Roth	False Sowthistle	Flowers	Asteraceae
8.	<i>Rhaphonticum acaule</i> (L.) DC.	Coffee plum	Flowers	Asteraceae
9.	<i>Scorzonera undulate</i> Vahl	Black Salsify	Aerial parts	Asteraceae
10.	<i>Tagetes minuta</i> Linn.	Wild Marigold	Flowers	Asteraceae

**Table 2:** Summary of Extractives Formulations

Concentration (%)	Amount of Stock Solution (ml)	Amount of Benzene (ml)	Amount of Emulsifiable Water (ml)	Total Amount (ml)
0.25	2.50	22.50	475.00	500.00
0.50	5.00	20.00	475.00	500.00
1.00	10.00	15.00	475.00	500.00
1.50	15.00	10.00	475.00	500.00
2.00	20.00	5.00	475.00	500.00

#### 4. Phago-Repellent Bioassay

For testing the repellent effect of asteraceous extracts okra leaves were used as food for *E. vittella* larvae treated with different concentrations. The treated foods were kept in jar (23cm x 10 cm) on moist filter paper. Thirty (15 male and 15 female), 24 hours starved *E. vittella* larvae were released in each jar along with control. The treated seeds were dipped in Benzene + emulsified water only. After four hours of the release *E. vittella* larvae. Treated *E. vittella* larvae either repelled and forced them to move from treated jars A to an empty jar B through the plastic pipe. The ones found in plastic pipe were considered repelled individuals. The repellency data (in treated and untreated jars) and alive (in empty or untreated jars) were recorded for 6hr at an interval of 6 hours for each observation (Sighamony *et al.* 1984, Tripathi *et al.* 2000, 2001, Trivedi and Chandel, 2009) [24, 25, 26, 27]. The data was collected on the number of *E. vittella* larvae which reached the treated food and repellency over control was recorded and calculated using Abbott formula (Abbott, 1925) [28].

#### 5. Result and Discussion

The data depicted from the (table 1 and figure 1, 2, and 3) results in the present experiment of repellent test against larvae of *Earias vittella*. All the plant extracts showed a good Phago-repellent property, when compared with the control. The repellent property was observed based on their EC<sub>50</sub> values as: *C. intybus* (0.0823) > *I. racemosa* (0.1043) > *T. minuta* (0.1620) > *C. cinerariaefolium* (0.1677) > *M. duriaeri* (0.1788) > *S. undulate* (0.3127) > *A. paniculata*, (0.3805) > *R. acaule* (0.4405) > *R. tingitana* (0.6368) > *C. odorata* (0.8870), respectively. The relative EC<sub>50</sub> values of the above extracts are in descending order, 10.7700 > 8.5043 > 5.4753 > 5.0227 > 4.9608 > 2.8365 > 2.3311 > 1.3929 > 1.0000 times as repellents, where as *C. odorata* is taken as a unit.

In the support of present finding the various asteraceous extractives has been reported by several.

Entomologist among them effect of *Cichorium intybus*, *Inula racemosa*, *Tagetes minuta* and *Mantisalca duriaeri* extracts against pulse beetle, *Callosobruchus chinensis* Linn. (Bhartii and Chandel 2017) [29]. in the same way a number of Entomologist worked on several indigenous plant extractives are tested for their repellent biopotency and exhibited strong biopotency against pulse beetle, *Callosobruchus maculatus* (Maredia *et al.* 1992, Rahman and Talukder 2006 and Tripathi and Upadhyay, 2009) [30, 31, 32].


Similarly, different formulation *Cleome gynandra* extractives gave significant repellent efficacy to spider mite (*Tetranychus urticae* Koch). *Ocimum suave* was found to repel the tick *Rhipicephalus appendiculatus* (Esther *et al.* 1995) [33]. Aqueous and methanolic plant extracts of *T. minuta*, which exhibited high repellency (IR = 0.04) against repellent effectiveness against the red flour beetle, *Tribolium castaneum* (Herbst) (Coleoptera Tenebrionidae). The application of these botanicals may be promising in protecting of stored grains against coleopteran pests. (Padin *et al.* 2013) [34]. *Alpinia galanga* (L.) species of termites, *Coptotermes gestroi* (Wasmann) and *Coptotermes curvignathus* (Holmgren) (Isoptera: Rhinotermitidae). Repellent activity shows that 250 ppm of 1,8-cineol caused 50.00 ± 4.47% repellency for *C. gestroi* (Sagheer *et al.* 2011, Salari *et al.* 2011, Khan *et al.* 2013, *et al.* 2011, Chandel and Singh 2016, Fauziah *et al.* 2015, Chandel and Singh, 2017) [35, 36, 37, 38, 39, 40]. The maximum repellent activity was observed at 500 ppm in methanol extracts of *N. nucifera*, ethyl acetate and methanol extract of *P. nigrum* and methanol extract of *T. ammi* and the mean complete protection time ranged from 30 to 150 min with the different extracts tested. (Liu *et al.* 2013) [41].

**Table 3:** Calculation of Log Conc./Probit Repellency Regression column of asteraceous extractives against *Earias vittella* larvae


Plant Extracts	Het.*	X <sup>2</sup>	Regression Equation	Ec <sub>50</sub>	Fiducial Limit	Relative Ec <sub>50</sub>
<i>Acmella paniculata</i>	3	0.96	Y=0.72X +3.93	0.3805	M1=2.9870 M2=1.1284	2.836
<i>Chromolaena odorata</i>	3	1.88	Y=0.90X+3.21	0.8870	M1=2.3866 M2=0.1286	1.000
<i>C. cinerariaefolium</i>	3	1.59	Y=1.08X+2.66	0.1677	M1=2.3362 M2=1.9781	5.022
<i>Cichorium intybus</i>	3	0.28	Y=0.75X +4.1	0.0823	M1=2.1368 M2=1.278	10.770
<i>Inula racemosa</i>	3	4.02	Y=1.07X +2.70	0.1043	M1=2.3256 M2=1.9392	8.504
<i>Manisalca duriaeri</i>	3	0.52	Y=0.88X +4.42	0.1788	M1=1.8678 M2=0.1251	5.022
<i>Reichardia tingitana</i>	3	0.52	Y=0.88X +4.42	0.6368	M1=1.8678 M2=0.1251	1.392
<i>Rhaponticum acaule</i>	3	1.95	Y=0.88X +4.37	0.4405	M1=1.5188 M2=0.0148	2.331
<i>Scorzonera undulate</i>	3	0.20	Y=0.84X +4.50	0.3127	M1=1.9418 M2=0.0432	4.960
<i>Tagetes minuta</i>	3	0.76	Y=1.07X +3.03	0.1620	M1=2.0527 M2=1.6572	5.475

In case of X<sup>2</sup> was found non significant heterogeneous at P=0.05, Y=Probit repellency, X=Log Concentration X 10,


D.F. = Degree of Freedom, E.C.<sub>50</sub> = Concentration Calculated at given 50% repellency.



**Fig 1:** Calculation of log conc./Probit Repellency Regression column of extractives against *E. vittella* larvae based on EC<sub>50</sub>



**Fig 2:** Calculation of log conc./Probit Repellency Regression column of extractives against *E. vittella* larvae based on relative EC<sub>50</sub>



**Fig 3:** Log concentration probit repellency regression column of comparative EC<sub>50</sub> and Relative EC<sub>50</sub> values of extracts against *E. vittella* larvae

## 6. Conclusion

Conclusively, the present investigation revealed that there appears prospects in selected botanicals *Cichorium intybus*, *Inula racemosa* and *Tagetes minuta* was registered promising repellency with minimum EC<sub>50</sub> values to the *E.vittella* larvae, when compared to other extracts. Among them *Cichorium intybus* elicited a strong defense response to counteract The growth and development of *E.vittella* larvae was significantly retarded when they fed on *Cichorium intybus* followed by *Inula racemosa* and *Tagetes minuta* extracts as compared to *C. odorata* treated okra leaves. The data depicted in table 1 and figure 1 indicated that result based on their relative EC<sub>50</sub> values the extracts of *Cichorium intybus* showed highest relative repellency (7.41) to the early emerged treated beetle followed by *Inula racemosa* (6.39), *Tagetes minuta* (5.88) and *Chrysanthemum cinerariaefolium* (5.42) times more repellent than *C. odorata* (1.00), which is taken as unit. The above selected plant materials can be used in protection of okra plant from *E.vittella* larval infestations under laboratory conditions. Thus, *C. intybus* and *Inula racemosa* aerial parts extract significantly reduced the population and damage caused by the major insect pests of okra recorded at Kanpur, India, compared to the untreated control. The *C. intybus*, *Inula racemosa* and *T.minuta* aerial parts extracts can be used effectively by farmers as a component of integrated management of pests of okra in India. This increase might be attributed to the plant's strong repellent properties. Importantly, selected extract from *C.oderata* leaves were also able to retard larval infestation of *E.vittella*. We conclude that defense metabolites from *C. intybus* and *Inula racemosa* possess strong repellent activity against *Plutella xylostella* (Linn.) even at lower concentrations revealed by present study and corroborated by earlier reports (Manish Kumar, 2017) [42].

## 7. Acknowledgements

Authors would like thank to Principal D.B.S. College, Kanpur, Uttar Pradesh for providing the necessary facilities. Prof. N.D. Pandey, Retd., Head, Division of Entomology, C S Azad University of Agriculture and Technology, Kanpur for rendering their support and help for the completion of this manuscript.

## 8. References

1. Radake SG, Undirwade RS. Seasonal abundance and

insecticidal control of shoot and fruit borer, *Earias* spp. on okra, *Abelmoschus esculentus* (L.). Indian J Entomol 1981;43:283-287.

- Kumar DS, Tony DE, Kumar AP, Kumar KA, Rao DBS, Nadendla RA. Review On: *Abelmoschus esculentus* (Okra), International Research Journal Phramceutical Applied Science 2013;3(4):129-132.
- Konar A, Rai L. Efficacy of some insecticides against shoot and fruit borer *Earias vittella* Fab. and *Earias insulana* (Boisd.) of okra (*Abelmoschus esculentus* L. Moench). Environ. Ecol 1990;8(1):410-413.
- Chaudhary HR, Dadheech LN. Incidence of insects attacking okra and the avoidable losses caused by them. Ann. Arid Zone 1989;28(3-4):305-307.
- Park HJ, Baek MY, Cho JG, Seo KH, Lee GY, Moon SJ, et al. Insecticidal alkaloids on aphids from *Corydalis turtschaninovii* tubers. J Korean Soc. Appl. Biol. Chem 2011;4:345-352.
- Chandel BS, Tripathi S, Singh A. Laboratory bioassay of N- (P-methoxy Phenyl-2 hydroxy salicylamide and N- (p-chlorophenyl) -2 hydroxysuccinimide as repellent against pulse beetle, *Callosobruchus chinensis* L. (Coleoptera: Bruchidae). International Journal of Entomology Research 2018;3(1):161-266.
- Dubey A, Gupta R, Chandel BS. Efficacy of *Acorus calamus*, *Vitex negundo* and *Ageratum conyzoides* against tobacco caterpillar *Spilarctia obliqua* Walker. Indian. J Entomology 2004;66(3):238-240.
- Chandel BS, Arti Singh. Repellent biopotency of *Cichorium intybus*, *Inula racemosa*, *Tagetes minuta* and *Mantisalca duriaeri* Aquash formulations against pulse beetle, *Callosobruchus chinensis* Linn. (Coleoptera: Bruchidae). Life Science Bulletin 2016;13(2):65-68.
- Dang QL, Lee GY, Choi YH, Choi GJ, Jang KS, Park MS, et al. Insecticidal activities of crude extracts and phospholipids from *Chenopodium ficifolium* against melon and cotton aphid, *Aphis gossypii*. Crop Prot 2010;29:1124-1129.
- Antonius AB, Hagazy G. Feeding deterrent activity of certain botanical extracts against cotton leafworm, *Spodoptera littoralis* Baised. Am. Agric. Sci 1987;32(1):719-729.
- Ahmed KS, Itino T, Ichikawa T. Effects of plant oils on oviposition preference and larval survivorship of

- Callosobruchus chinensis* (Coleoptera: Bruchidae) on azuki bean. Applied Entomology and Zoology 1999;34(4):547-550.
12. Al-Lavati HT, Azam KM, Deadman ML. Insecticidal and repellent properties of subtropical plant extracts against pulse beetle, *Callosobruchus chinensis*. Sultan Qaboos Univ. J Sci. Res. Agric. Sci 2002;7:37-45.
  13. Schmidt GH, Streloke M. Effect of *Acorus calamus* Linn. (Araceae) oil and its main compound B- Asarone on *Prostephanus truncatus* Horn (Coleoptera: Bostrichidae). J Stored Product Research 1994;30(3):227-235.
  14. Pavela R. Insecticidal activity of essential oils against cabbage aphid *Brevicoryne brassicae*. J Essent. Oil Bearing Plants 2006;9:99-106.
  15. Rashid T, Chen J, Mcledo P. Toxicity of newly isolated piperidine alkaloids from the red imported fire ant, *Solenopsis invicta*, against the green peach aphid, *Myzus persicae* (Sulzer). Adv. Entomol 2013;1:20-23.
  16. Srivastava AS, Awasthi GP. An insecticide from the extract of the plant *Adhatoda vasica* harmless to man. P roc. 10<sup>th</sup> Ent. Congr. Exp 1958;2:245-246.
  17. Mendes Queiroz RO, Zani CL. Screening of asteraceae (Compositae) plant extracts for larvicidal activity against *Aedes fluviatilis* (Diptera: Culicidae). Mem. Inst. Oswaldo Cruz 1997;92:565-570.
  18. Kim SK, Kim YC, Lee S, Kim JC, Yun MY, Kim IS. Insecticidal activity of rhamnolipid isolated from pseudomonas sp. EP-3 against green peach aphid (*Myzus persicae*). J Agric Food Chem 2011;59(3):934-8.
  19. Jilani G, Saxena RC, Rueda BP. Repellency and toxicity of some plant oils and their terpene components to *Sitotroga cerealella* Oliver (Lipidoptera, Gelechiidae), Tropical Science 1985;25:249-252.
  20. Abubakar MS, Abdurahman EM, Haruna AK. The repellent and antifeedant properties of *Cyperus articulatus* against *T. castaneum*. Phytotherapy Research 2000;14:281-283.
  21. Ramamurthy R, Rajaram V, Rajendran LM. Efficacy of plant products against the storage pest bruchides in black gram. Legume Research 2002;32:34.
  22. Kamaraj C, Rahuman AA, Bagavan A, Gandhi E, Abdul Abduz Zahir, Thirunavukkarasu AA *et al.* Larvicidal and repellent activity of medicinal plant extracts from Eastern Ghats of South India against malaria and filariasis vectors. Asian Pacific Journal of Tropical Medicine 2011;4(9):698-705.
  23. Chandel BS, Sengar HS. Insectifuge effect of neem, *Azadirachta indica* A. Juss and their derivatives against sawfly, *Athalia lugens proxima* Klug. (Hymenoptera: Tenthredinidae). International Journal of Zoology Studies 2018;3(1):53-59.
  24. Sighamony S, Anees I, Chandrakala TS, Osmani Z. Natural products as repellents for *Tribolium castaneum* Herbst. International Pest Control 1984, 156-157.
  25. Tripathi AK, Prajapati V, Agarwal KK, Khanuja SPS, Kumar S. Repellency and toxicity of oil from *Artimesia annua* to certain stored product beetles. Jour. Econ. Ent 2000;93(1):43-47.
  26. Tripathi MK, Sahoo P, Das BC, Mohanti S. Efficacy of botanical oils, plant powders and extracts against *Callosobruchus chinensis* Linn. Attacking black gram (CV. T9). Legume Res 2001;24:82-86.
  27. Trivedi SS, Chandel BS. Repellent Efficacy of *Adhatoda vasica*, *Pongamia glabra* and *Momordica charantia* plant extract against *Spilarctia obliqua* Walker (Lepidoptera: Arctiidae). International Journal of Biotechnology and Biochemistry 2009;5(4):469-467.
  28. Abbott WS. A method of computing the effectiveness of insecticides. Journal of Economic Entomology 1925;18:265-267.
  29. Bharti A, Chandel BS. Insectifuge Biopotency of *Chrysanthemum cinerariifolium*, *Cichorium intybus*, *Inula racemosa* and *Tagetes minuta* extractives against okra aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae). International Journal of Fauna and Biological Studies 2017;4(3):133-138.
  30. Maredia KM, Segura OL, Mihm JA. Effects of neem, *Azadirachta indica*, on six species of maize insect pests. Tropical Pest Management 1992;38(2):190-195.
  31. Rahman A, Talukder FA. Bioefficacy of some plant derivatives that protect grain against the pulse beetle, *Callosobruchus maculatus*. J Insect Science 2006;6:11-14.
  32. Tripathi AK, Upadhyay S. Repellent and insecticidal activities of *Hyptis suaveolens* leaf essential oil against four stored grain coleopteran pests Journal of Stored Prod. Res 2009;40:317-330.
  33. Esther Mwangi N, Ahmed Hassanali, Suliman Essuman, Edward Myandat, Lambert Moreka, Mark Kimondo. Repellent and acaricidal properties of *Ocimum suave* against *Rhipicephalus appendiculatus* ticks. Experimental & Applied Acarology 1995;19(1):11-18.
  34. Padin SB, Fuse C, Urrutia MI, Dal Bello GM. Toxicity and repellency of nine medicinal plants against *Tribolium castaneum* in stored wheat, Bulletin of Insectology 2013;66(1):45-49.
  35. Sagheer M, Mansoor-ul-Hasan, Latif MA, Iqbal J. Evaluation of some indigenous medicinal plants as a source of toxicant, repellent and growth inhibitors against *Tribolium castaneum* (Coleoptera: Tenebrionidae). Pak. Entomol 2011;33:87-91.
  36. Salari E, Ahmadi K, Dehyaghobi RZ, Purhematy A, Takaloozadeh HM. Toxic and repellent effect of harmful (*Peganum harmala* L.) acetonetic extract on several aphids and *Tribolium castaneum* (Herbst). Chil. J agric. Res 2011;72:147-151.
  37. Khan FZA, Sagheer M, Mansoor-ul-Hasan, Saeed S, Ali K, Gul HT, *et al.* Toxicological and repellent potential of some plant extracts against stored product insect pest, *Tribolium castaneum* (Herbst.) (Coleoptera: Tenebrionidae). Int. J Biosci 2013;3:280-286.
  38. Chandel BS, Singh A. Repellent biopotency of *Cichorium intybus*, *Inula racemosa*, *Tagetes minuta* and *Mantisalca duriaeri* Aquash formulations against pulse beetle, *Callosobruchus chinensis* Linn. (Coleoptera: Bruchidae). Life Science Bulletin 2016;13(2):65-68.
  39. Fauziah Abdullah, Partiban Subramanian, Halijah Ibrahim, Sri Nurestri Abdul Malek, Guan Serm Lee, Sok Lai Hong. Chemical composition, antifeedant, repellent, and toxicity activities of the rhizomes of galangal, *Alpinia galanga* against Asian subterranean Termites, *Coptotermes gestroi* and *Coptotermes curvignathus* (Isoptera: Rhinotermitidae). J Insect Sci 2015;15(1):112-118.
  40. Chandel BS, Singh A. Entomotoxicity of *Chromolaena odorata*, *Tagetes minuta* and *Reichardia tingitana* in

suppressing oviposition and adult emergence of *Callosobruchus chinensis* (L.) infesting stored chickpea seeds in U.P. International Journal of Zoology Studies 2017;2(6):38-44.

41. Liu CH, Mishra AK, Tan RX. Repellent, insecticidal and phytotoxic activities of isoalantolactone from *Inula racemosa*. Crop Protection 2006;25:508-511.
42. Manish Kumar, Tripathi RC, Chandel BS. Potential impact of *Inula racemosa*, *Cichorium intybus* and *Mantisalca duriaeri* against *Plutella xylostella* Linn. (Lepidoptera: Noctuidae). Life Science Bulletin 2017;14(1):65-69.