



E-ISSN 2347-2677

P-ISSN 2394-0522

[www.faujournal.com](http://www.faujournal.com)

IJFBS 2022; 9(4): 22-27

Received: 23-05-2022

Accepted: 27-06-2022

**Anju Bharti**

Biopesticide Lab, Department of  
Zoology, D.B.S. College, Kanpur,  
Uttar Pradesh, India

**Alok Kumar Srivastava**

Biopesticide Lab, Department of  
Zoology, D.B.S. College, Kanpur,  
Uttar Pradesh, India

**BS Chandel**

Biopesticide Lab, Department of  
Zoology, D.B.S. College, Kanpur,  
Uttar Pradesh, India

## Field assessment of biorational naturally occurring botanicals for management of tobacco caterpillar, *Spilarctia obliqua* Walker (Lepidoptera: Arctiidae) on cabbage, *Brassica oleracea* var. *capitata*

Anju Bharti, Alok Kumar Srivastava and BS Chandel

### Abstract

Pesticides have been a major contributor to the growth of agricultural productivity and food supply. Pesticides were a key factor in significant agricultural productivity growth during the last century and continue to be a critical factor in reducing crop damage. An experiment was carried out to test the insecticidal efficacy of five plant extracts against third instars larvae of *Spilarctia obliqua* Walker under the field conditions. The alcoholic extracts of leaves of pavettia, *Adhatoda vasica* Nees (Acanthaceae), black nightshade, *Solanum nigrum* Linn. (Solanaceae), arial parts of ariapple, *Lantana camara* Linn. (Verbanaceae), seeds of hulhul, *Cleome monophylla* Linn. (Capparidaceae) and rhizome of *Zingiber officinalis* Linn. (Zingiberaceae) were tested on cabbage, *Brassica oleracea* var. *capitata*. Among them 2.0% extracts showed mortality to the third instars larvae of *Spilarctia obliqua* Walk. Among them, *Adhatoda vasica* leaves extract showed highest larval mortality. Based on mean mortality per cent findings are arranged in the following descending order of merit as: *A. vasica* (66.35) > *C. monophylla* (64.17) > *L. camara* (62.62) > *S. nigrum* (62.13) > *Z. officinalis* (59.64) > Control (06.14), respectively. All the extracts were considerably superior over control. The study indicated that *Adhatoda vasica* leaves extract was the most suitable for cabbage insect-pest management from the severe infestation of *Spilarctia obliqua*. These botanicals can be recommended for the farmer's use to control *Spilarctia obliqua* larvae.

**Keywords:** Cabbage, *brassica oleracea* var. *capitata*, *Spilarctia obliqua*, *Adhatoda vasica*, *Cleome monophylla*, *Lantana camara*

### 1. Introduction

Bihar hairy caterpillar, *Spilarctia* (*Dicrisia* or *Spilosoma*) *obliqua* Walker (Arctiidae: Lepidoptera) is highly polyphagous and cosmopolitan insect pest of sporadic nature in India (Kabir and Khan, 1968; Steven *et al.* 2007<sup>[1, 2]</sup>, it attacks vegetables, pulses, oilseeds, cereals, mulberry, medicinal, aromatic plants and bast fibre crops (Bhattacharya *et al.* 1995)<sup>[3]</sup>.

The infestation caused by *Spilarctia obliqua* various vegetables like Cabbage, *Brassicra oleraea* var. *capitata*, Broccoli, *Brassicra oleraea* var. *italica*, Cauliflower, *Brassicra oleracea* var. *botrytis* and jute, *Corchorus olitorius*, *C. capsularis*, mustard, *Brassicra juncea*, sunnhemp, *Crotalaria juncea*, fiber, oil yielding crop and often causes severe qualitative and quantitative damage (Varatharajan *et al.* 1998, Singh and Varatharajan, 1999, Begum *et al.* 2000, Awmack and Leather, 2002, Dhingra *et al.* 2000, Gotyal *et al.* 2013 and<sup>[4, 5, 6, 7, 8, 9]</sup>.

Cabbage, *Brassicra oleraea* var. *capitata* contains bio-chemicals that are thought to help prevent cancer. Cabbage might change the way estrogen is used in the body, which might reduce the risk of breast cancer. Cabbage is packed with nutrients, excellent source of vitamin K and vitamin C. it helps improve digestion and may lower blood pressure may help keep your heart healthy, could help lower cholesterol levels (Rahman *et al.* 2020)<sup>[10]</sup>.

There are several constraints in increasing the productivity of vegetables of which the losses due to insect pests is one of the major concern (Das and Chaudhuri, 2005)<sup>[11]</sup>. Among the insect, larvae of *S. obliqua* is one of the major which causing substantial yield loss as well as vegetable quality. (Roy Barik 2013)<sup>[12]</sup>.

Management of this pest in due time is very important as delay may lead to even complete defoliation of vegetables and brassicaceous crop. Although many insecticides are effective against this pest, chemical control is unfit for its control.

**Corresponding Author:****Anju Bharti**

Biopesticide Lab, Department of  
Zoology, D.B.S. College, Kanpur,  
Uttar Pradesh, India

Excessive use of synthetic insecticides has led to problems has been raised about the ill effects to human health like hazardousness, un-biodegradability and relative resistance of *S. obliqua*, affecting environment and increase the casts of insect control (Mandal and Bhattacharya 2003 <sup>[13]</sup>).

Host plant availability and quality is a key determinant of herbivorous insect population dynamics and development of insect and adult performance (Gautam *et al.* 2003) <sup>[14]</sup>. A large number of insect-pests as, cabbage butterfly, *Pieris brassicae* Linn., tobacco caterpillar, *Spodoptera litura* Fabr., Bihar hairy caterpillar, *Spilarctia obliqua* Walk., cabbage borer, *Hellula undalis* Fabr., mustard sawfly, *Athalia proxima* Klug., cabbage semilooper, *Trichoplusia ni* Hub. and cabbage leaf webber, *Crociodolomia binotalis* Zell. are limiting factors of crops (Kulat *et al.* 1999, Bajpai and Chandel, 2010,) <sup>[15, 16]</sup>. The pest is distractive in its larval stages. The damaged plant shows stunty growth and results in deterioration of yield (Chandel and Singh, 2017 and Bharti *et al.* 2021) <sup>[17, 18]</sup>.

Increasing attention in environmental safety has triggered interest in pest control approaches through eco-friendly plant-based pesticides. Botanical pesticidal constituents are effective against *Spilarctia obliqua* destructive pests of cabbage, *Brassicae oleraceae* var. *capitata*. More importantly, they are widely available, inexpensive, accessible, rapidly

biodegradable, and have little toxicity to beneficiary agents. Therefore present investigation was carried out to develop eco-friendly, non-hazardous, non- residues and biorational extractives as new alternative aspect of synthetic chemical for the control of *Spilarctia obliqua* on cabbage.

## 2. Material and Methods

### 2.1 Survey and collection of Plant Materials

In the present investigation, the efficacy of five plant origin insecticide i.e. leaves of pavettia, *Adhatoda vasica* Nees (Acanthaceae), black nightshade, *Solanum nigrum* Linn. (Solanaceae), arial parts of ariapple, *Lantana camara* Linn. (Verbanaceae), seeds of hulhul, *Cleome monophylla* Linn. (Capparidaceae) and rhizome of *Zingiber officinalis* Linn. (Zingiberaceae) were collected from vicinity of Kanpur.

### 2.2 Extraction of plant materials

In the present investigation alcoholic extract of leaves of *Adhaloda vasica* Nees, *Solanum nigrum* Linn, seeds of *Cleome monophylla* Linn, Rhizome of *Zingiber officinatis* Rose, Arial parts of *Lantana camara* Linn. were dried in shade make them powder, extracted them with the help of the soxhlet apparatus under the field conditions (Bharti and chandel, 2017) <sup>[19]</sup>.

**Table 1:** List of indigenous plant materials and their details

Scientific Name	Vernacular Name	Part Used	Family
<i>Adhatoda vasica</i> Nees	Pavettia	Leaves	Acanthaceae
<i>Cleome monophylla</i> Linn.	Hulhul	Seeds	Capparidaceae
<i>Lantana camara</i> Linn.	Ariapple	Arial parts	Verbanaceae
<i>Solanum nigrum</i> Linn.	Black nightshade	Leaves	Solanaceae
<i>Zingiber officinalis</i> Linn.	Ginger	Rhizome	Zingiberaceae

### 2.3 Preparation of 50 percent stock solution from pure extract

50ml. Extract in each case was taken into reagent bottle and 50ml. Benzene was added in it to dissolve the constituents of the materials. This was the 50 percent stock solution, the mouth of the bottles were stopped with airtight corks and kept in refrigerator.

### 2.4 Formulations of Different concentrations

The different concentrations of the insecticides were prepared from the stock solution using benzene as solvent and Triton X-100 as emulsifier. The level of solvents and emulsifier were kept constant at the rate of 5 per cent and 0.5 per cent, respectively, in the final spray.

### 2.5 Preparation of 0.5 Per cent Emulsifiable Water

0.5 ml. of Triton X-100 was accurately measured into a large bottle with the help of a measuring cylinder, then of distilled water was added and bottle was shaken well to dissolve the

emulsifier. Thus emulsifiable water of 0.5 per cent strength was obtained and used for the preparation of different concentrations of the extracted materials (Schmidt and El, 1997) <sup>[20]</sup>.

### 2.6 Preparation of Various Concentrations

To make various concentration of extract the required quantity of the stock solution was calculated with the help of following formula:

$$\text{Amount of Stock Solution} = \frac{\text{Amount required} \times \text{Concentration required}}{\text{Concentration of Stock Solution}}$$

The calculated amount of various ingredients required to make different concentrations from the 50 per cent stock solution and amount of ingredients taken are presented in the following table:

**Table 3:** Preparation of different formulations of the selected plant materials

Concentration (%)	Amount of Stock Solution (ml)	Amount of Benzene (ml)	Amount of Emulsifiable Water (ml)	Total Amount (ml)
0.50	5.00	20.00	475.00	500.00
1.00	10.00	15.00	475.00	00.00
2.00	20.00	5.00	475.00	500.00

### 3. Experimental Procedure

#### The Toxicity Test of soxhlet extract under field condition Insecticidal Test

A preliminary trial was carried out to test the insecticidal properties of five plant extracts in three concentrations in the insectary field at fattepur village affiliated to Department of Zoology, D.B.S. College, C.S.J.M. University, Kanpur. All the three concentrations of extract were sprayed on Cabbage, *Brassica oleracea* var. *capitata* plants. After treatment, the number of larvae remained in each plot were recorded at an interval of 24 hrs. 48 hrs. And 72 hrs. The percentage reduction of larvae was calculated and the percentage reduction of larvae was converted into angular values (sign/percentage). The data were statistically analysed (Finny, 1938) [21].

The concentrations of 0.5, 1.0 and 2.0% of all the thirteen plant extracts were prepared in water using triton @ 0.5% as emulsifier and benzene @ 5.0% as solvent. The laboratory experiments and field trials to test the insecticidal effectiveness were carried out with three replications during November and December 2020 in the laboratory and field of D.B.S.P.G. College, Kanpur (India).

In laboratory, the plant extracts were tested by dry film technique. Ten plant extracts and their three concentrations (0.5, 1.0 and 2.0 per cent) including untreated control was replicated thrice. Paired petri-dishes (10 cm diameter) were selected for this insecticidal test. For preparing a film 1.0 ml of extract preparation was poured into a petri-dish and it was

gently shaken under an electric fan till the liquid phase evaporated leaving behind on the glass surface an uniform dry film of plant extract. Ten 24 hrs starved third instars larvae were introduced into each paired petri-dish. The larvae were given a continuous exposure to the plant extract films for 2 hrs. After the treatment the larvae from each paired petri-dishes were transferred to separate clean jar (23 cm X 15 cm) containing fresh okra and mustard leaves as food. Mouth of jar was kept covered by a piece of muslin cloth kept in position with rubber band around it. Observations on the mortality of larvae were recorded after 24, 48 and 72 hrs of their release.


Similarly, the field experiment was conducted in randomized block design (RBD) having plot size of 1.0 m X 2.0 m. Care was taken to protect the fresh larvae coming in the plot from outside. *Acorus calamus*, *Vitex negundo*, *Ageratum conyzoides* and other remaining seven plant extracts of 0.5%, 1.0% and 2.0% concentrations with three replications were sprayed over the okra and mustard plant and ten larvae were released in each plot. The spray was applied with an atomizer. Observation on the mortality of larvae of *S. obliqua* was recorded after 24, 48 and 72 hrs on their release.

The data on percent infestation was subjected to ANOVA after angular transformation. Modified Abbot's formula was used for calculation of percentage reduction of pest population over control (Abbott, 1925 and Gomez and Gomez, 1984) [22, 23].


**Table 1:** Mean mortality % of *Spilarcia obliqua* in different exposure period irrespective of concentrations under field conditions

Treatment	Mean Mortality percent after						Mean	%
	24 hrs.		48 hrs.		72 Hrs.			
	T <sub>1</sub>	TBV <sub>1</sub>	T <sub>2</sub>	T.B.V. <sub>2</sub>	T <sub>3</sub>	T.B.V. <sub>3</sub>	G.T.	TBV
<i>A. vasica</i> Nees	61.18	76.6	68.23	86.2	69.64	87.9	66.35	83.9
<i>C. monophylla</i> Linn.	59.93	74.9	64.57	81.6	68.00	86.0	64.17	81.0
<i>L. camara</i> Linn.	56.85	70.1	59.80	74.1	71.21	89.6	62.62	78.8
<i>S. nigrum</i> Linn.	51.74	61.6	61.14	76.7	73.52	91.9	62.13	78.1
<i>Z. officinalis</i> Linn.	54.49	66.3	60.54	75.7	63.90	80.7	59.64	74.4
Control (Untreated)	00.00	0.0	00.00	0.0	10.00	18.44	6.14	1.1


TBV=Transform Back Value, T=Treatment, G.T= Grant Total




**Fig 1:** Mean mortality % of *S. obliqua* after 24 hrs. Under field conditions



**Fig 2:** Mean mortality % of *S. obliqua* after 48 hrs. Under field conditions



**Fig 3:** Mean mortality % of *S. obliqua* after 72 hrs. Under field conditions



**Fig 4:** Over all mean mortality % of *S. obliqua* under field conditions

**4. Result and Discussion**

Bioassay studies using third instars larva of *S. obliqua* for calculating median Effective concentration (EC) for all the

test 50 insecticides were as follows:

Third instars larvae of *Spilarctia obliqua* Walker under the field conditions. The alcoholic extracts of leaves of pavettia,

*Adhatoda vasica* Nees (Acanthaceae), black nightshade, *Solanum nigrum* Linn. (Solanaceae), arial parts of ariapple, *Lantana camara* Linn. (Verbanaceae), seeds of hulhul, *Cleome monophylla* Linn. (Capparidaceae) and rhizome of *Zingiber officinalis* Linn. (Zingiberaceae) were tested on cabbage, *Brassica oleraea* var. *capitata*. Among them 2.0% extracts showed mortality to the third instars larvae of *Spilarctia obliqua* Walk. Among them, *Adhatoda vasica* leaves extract showed highest larval mortality.

Data depicted from findings that alcoholic extracts of leaves of pavettia, *Adhatoda vasica* Nees (Acanthaceae), showed highest larval mortality and placed on the top of merit gave 66.35 per cent mean mortality followed by seeds of hulhul, *Cleome monophylla* Linn. (Capparidaceae) (64.17), arial parts of ariapple, *Lantana camara* Linn. (Verbanaceae), (62.62), black nightshade, *Solanum nigrum* Linn. (Solanaceae), (62.13), rhizome of *Zingiber officinalis* Linn. (Zingiberaceae) (59.64) and Control (06.14), respectively.

In the support of above findings many entomologist like Talukder, 1989, Tohnishi *et al.* 2005, Nair *et al.* 2007, Gao *et al.* 2008, Sonu *et al.* 2008, Polan *et al.* 2009, Salim *et al.* 2015 used *Adhatoda vasica* and other botanicals against number of insect pest particularly *Spilactia obliqua* [24, 25, 26, 27, 28, 29, 30].

Sadek 2003 used crude methanolic extracts of *Adhatoda vasica* leaves on the feeding of *Spodoptera littoralis* larvae was investigated in the laboratory. Feeding on fresh leaves resulted in 100% mortality of larvae after 26 days of unsubstantial growth. The extract exhibited strong antifeedant and toxic activity against the larvae when applied either on leaf discs or incorporated into artificial diet [32].

Sawadogo *et al.* 2009 reported that *Cleome viscosa* L. extract showed significant insecticidal activity against cowpea pod sucking bugs, *Clavigralla tomentosicollis*.

Borooah 2011 reported that *Adhatoda vasica* extract exhibited anti insect activity and extensively used in pest control.

Ngeggha, *et al.* 2022 stated that pest management is being confronted with immense economic and environmental issues worldwide because of massive utilization and over-reliance on pesticides. The non-target toxicity, residual consequence, and challenging biodegradability of these synthetic pesticides have become a serious concern, which urgently requires the alternative and prompt adoption of sustainable and cost-effective pest control measures [34].

Rajashekar *et al.* 2012 reported that oil of *L. camara* was significantly kill the stored grain pests and stored maize could be protected from the infestation of *S. zeamais* [35].

Therefore the present investigation showed ecifriendly, biorational and appropriate alternative of synthetic insecticide may provide platform for developing the appropriate alternative of synthetic insecticide for the control of *S. obliqua* infesting on cabbage, *Brassica oleraea* var. *capitata* under field trials.

## 5. Acknowledgement

The authors thank the Principal D.B.S. College, Kanpur and Head, Department of Zoology, D.B.S. College, Kanpur, India, for providing the necessary facilities to perform the present study. Prof. R.A. Tripathi, farmer, Head, Division of Entomology, C S Azad University of Agriculture and Technology, Kanpur for rendering their support and help for the completion of this work and manuscript.

## References

1. Kabir AKMF, Khan SA. Bioassay of some insecticides for the control of jute hairy caterpillar, *Diacrisia obliqua* Walk. Indian Journal of Agricultural Science. 1968;6(1-2):131-138.
2. Steven E, Lei SZ, Zilberman D. The Economics of Pesticides and Pest Control. International Review of Environmental and Resource Economics, 2007;1(3):271-326.
3. Bhattacharya PK, Ram WH, Sarker S, *Spilosoma obliqua* (Walker): economic importance, biology, host range and breeding for resistance in soybean. Agric. Rev. India. 1995;16(1, 2):23-40.
4. Varatharajan R, Singh AS, Keisa JT, Singh OD, Selvasundaram R. Life table of *Spilosoma obliqua* (Lepidoptera: Arctiidae) on sunflower. International J. Trop. Insect Sci. 1998;18:383-85.
5. Singh YR, Varatharajan R. Host range of Bihar hairy caterpillar, *Spilosoma obliqua* (Walker) (Arctiidae: Lepidoptera). Hexapoda. 1999;11:65-74.
6. Begum K, Rahman AKMM, Haque MA, Begum J, Uddin MM. Biology of jute hairy caterpillar, *Spilarctia obliqua* (Walker) on different food plants. Bangladesh J Ent. 2000;10:71-77.
7. Awmack CS, Leather SR. Host plant quality and fecundity in herbivorous insects. Ann. Rev. Ent. 2002;47:817-44.
8. Dhingra S, Bhandari JKS, Shankarganesh K. Relative resistance of Bihar hairy caterpillar to insecticide mixtures. J Ent. Res. 2007;31:209-12.
9. Gotyal BS, Satpathy S, Selvaraj K, Babu RV. Comparative biology of Bihar hairy caterpillar, *Spilosoma obliqua* on cultivated and wild jute species. Indian J Pl. Prot. 2013;41:219-21.
10. Rahman MS, Islam MN, Talukder FU, Sultan MT. Evaluation of insecticides for the management of jute hairy caterpillar, *Spilosoma obliqua* walk. (Lepidoptera: Arctiidae) in jute, *Corchorus Olitorius*, International Journal of Entomology Research. 2020;5(4):71-77.
11. Das M, Chaudhuri N. Life-fecundity table on incidence of *Spilosoma obliqua* on jute. Ann. Pl. Prot. Sci. 2005;13:319-23.
12. Roy N, Barik A. Influence of four host-plants on feeding, growth and reproduction of *Diacrisia casignetum* (Lepidoptera: Arctiidae). Entomol. Sci. 2013;16:112-18.
13. Mandal P, Bhattacharya AK. Azadirachtin, the most potent neem derivative: effect on *Spilosoma obliqua*. Indian J Ent. 2003;65(2):170-176.
14. Gautam Kaushal, Rao PB, Chauhan SVS. Insecticidal properties of some plants of family Asteraceae against *Spilosoma oliqua*. Indian J Ent. 2003;65(3):363-367.
15. Kulat SS, Nimbalkar SA, Radke SG, Tambe VJ. Evaluation of biopesticides and neem seed extract against *Helicoverpa armigera* on chickpea. Indian J Ent. 1999;61(1):19-21.
16. Bajpai Rachna, Chandel BS. Assessment of certain plant extracts as insecticides against cabbage butterfly, *Pieris brassicae* Linn. (Lepidoptera: Pieridae). International Journal of Biotechnology and Biochemistry. 2010;6(6):1003-1017.

17. Chandel BS, Singh A. Phagodeterrent bioefficacy of *Acorus calamus*, *Withania somnifera* and *Momordica charantia* against chickpea bruchids, *Callosobruchus chinensis* (Linn.) (Coleoptera: Bruchidae), Journal of Entomology and Zoology Studies. 2017;5(6):935-939.
18. Bharti Anju, Dubey Indrani, Chandel BS. Phago-repellent biopotency of *Cichorium intybus*, *Inula racemosa*, *Tagetes minuta* and *Chrysanthemum cinerariaefolium* as herbal alternatives to synthetic insecticides against *Earias vittella* Fabricius (Lepidoptera: Noctuidae) on Okra in Kanpur region. International Journal of Fauna and Biological Studies. 2021;8(3):11-17.
19. Bharti Anju, Chandel BS. Aphidicidal compatibility of certain ecofriendly asteraceous biorational extractives against okra aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae) under field conditions. International Journal of Fauna and Biological Studies. 2017;4(3):125-132.
20. Schmidt and El Evaporation, absorption and penetration of insecticidal ingredients of Indian essential *Acorus calamus* oil. Bollettino di Zoologia Agraria e di Bachicoltura. 1997;29(2):167-181.
21. Finney DJ. Probit analysis: a statistical treatment of the sigmoid responses in curve. Rev. ed. Cambridge University Press, London, 1938, 318.
22. Abbott WS. A method of computing the effectiveness of insecticides. Journal of Economic Entomology. 1925;18:265-267.
23. Gomez KA, Gomez AA. Statistical Procedures for Agriculture Research. John Weley and Sons. Inc., New York, 1984, 67-265.
24. Talukder, D., Khan, A.R., Hasan, M. Growth of *Diacrisia obliqua* (Lepidoptera: Arctiidae) with low doses of *Bacillus thuringiensis* Var. Kurstaki. Entomophaga. 1989;34(4):587-589.
25. Tohnishi M, Nakao H, Furuya T, Seo A, Kodama H, Tsubata K, et al. Flubendiamide a novel insecticide highly active against lepidopterous insect pests. J Pestic. Sci, 30, 354-360.
26. Nair N, Sekh K, Debnath M, Chakraborty S, Somchoudhury AK. Relative toxicity of some chemicals to bihar hairy caterpillar, *Spilarctia obliqua* Walker (Arctiidae, Lepidoptera). Journal of Crop Weed. 2007;3(1):1-2.
27. Gao H, Huang YN, Gao B, Li P, Inagaki C, Kawabata J. Inhibitory effect on  $\alpha$  glucosidase by *Adhatoda vasica* Nees. Food Chem. 2008;108:965-972.
28. Soni S, Anandjiwala S, Patel G, Rajani M. Validation of different methods of preparation of *Adhatoda vasica* leaf juice by quantification of total alkaloids and vasicine. Indian J Pharm. Sci. 2008;70:36-42.
29. Polan MS, Banu H, Islam MN, Haque SMA, Mosaddeque HQM. Field efficacy and evaluation of effective dose of some insecticides against jute hairy caterpillar, *Spilosoma obliqua* Walker. Bangladesh. J Jute Fib. Res. 2009;29(1-2):69-75.
30. Salim HA, Abed MS. Effect of botanical extracts, biological and chemical control against *Spilosoma obliqua* on cabbage (*Brassica oleracea*). Journal of Entomology and Zoology Studies. 2015;3(1):43-46.
31. Sadek M. Antifeedant and toxic activity of *Adhatoda vasica* leaf extract against *Spodoptera littoralis* (Lepidoptera: Noctuidae) Journal of Applied Entomology. 2003;127(7):396-404.
32. Sawadogo NMBAF, Dabire-Binso CL, Drabo I, Sanon A. Insecticidal Activity of Three Plants Extracts on the Cowpea Pod Sucking Bug, *Clavigralla tomentosicollis*, STÄL (Hemiptera: Coreidae). Pakistan Journal of Biological Sciences. 2009;12:1320-1324.
33. Borooah DD. Effect of *Adhatoda vasica* (L.) Neem leaf extract prepared by two different methods on mitosis of root meristematic cells of *Allium cepa* Linn. Journal of Advanced Laboratory Research in Biology. 2011;2(1):141-145.
34. Ngegha PM, Cui G, Khalid MZ, Zhong G. Use of Botanical Pesticides in Agriculture as an alternative to Synthetic Pesticides, 2022, Agriculture. 2022;12(5):600, 25-30.
35. Rajashekar Y, Ravindra KV, Bakthavatsalam N. Leaves of *Lantana camara* Linn. (Verbenaceae) as a potential insecticide for the management of three species of stored grain insect pests. Journal of Food Science and Technology -Mysore. 2012;51(11):85-89.