



International Journal of Fauna and Biological Studies

Available online at www.faunajournal.com

I
J
F
B
S

International
Journal of
Fauna And
Biological
Studies

E-ISSN 2347-2677

P-ISSN 2394-0522

Impact Factor (RJIF): 5.69

<https://www.faunajournal.com>

IJFBS 2025; 12(5): 106-110

Received: 15-08-2025

Accepted: 16-09-2025

Parna Bhadra

Department of Zoology, MES

Abasaheb Garware College

(Autonomous), Pune,

Maharashtra, India

Amol Salokhe

Department of Zoology, MES

Abasaheb Garware College

(Autonomous), Pune,

Maharashtra, India

A morphometric study of *Macrosiphum euphorbiae* (Thomas) (Hemiptera: Aphididae) on *Rosa* sp. from four different locations in Pune

Parna Bhadra and Amol Salokhe

DOI: <https://www.doi.org/10.22271/23940522.2025.v12.i5b.1134>

Abstract

Morphometric studies were carried out on four populations of *Macrosiphum euphorbiae* (Hemiptera: Aphididae) associated with four varieties of *Rosa* sp. Ten morphological characters were measured and the mean value of each character was compared to the same of the other three populations collected from different sites. Out of the ten characters (body length, body width, head width, siphunculus length, processus terminalis length, basal antennal segment), only length of siphunculus, antennae, body, width of body and ultimate rostral segment length indicated the occurrence of locally adapted populations. Morphometric analyses suggested not only about the occurrence of four morphologically differentiated groups within *M. euphorbiae* from four isolated localities but also a preference of undisturbed forest area against the urban or city area by the pest species. We have tried to study if aphids show response to genetic variation (rose varieties) in their host plant and local conditions.

Keywords: Aphid, host association, morphometry, local adaptations, taxonomic population

Introduction

Insects under the family Aphididae are serious and stubborn sapsucking pests under order Hemiptera infesting agricultural and horticultural crops [1]. They directly damage plants by sucking their nutrients and causing reduced vigour by pouring toxic saliva [2]. They damage the crops by making leaf and stem galls besides secreting honeydew in high amounts causing disturbance in stomatal activity by blocking it, thus, reducing photosynthesis because of fungal development that invites viral diseases [2]. They are unique in having complex lifecycles with alternation of generation, multiple host plants, telescopic generations, polymorphism, high fecundity and parthenogenetic viviparity [3]. Temperate regions of the world show higher diversity of aphids than tropics [4]. Around 800 species are known from India and 385 of these are endemic [5]. Plant feeding population of insects and their population dynamics is influenced by the genetic variation of plants [6, 7]. Insects show variable ability to survive on different host plants [8, 9] and also varieties of the same plant species expecting that these plants are showing genetic variation [10]. Two of the *Chaitophorus populicola* aphid populations among four on *Populus angustifolia* evidently showed that they were adapted to the local host plant, in the third the herbivore was maladapted to the local host-plant population while the fourth population had no evidence for adaptation [11].

This study deals with collections of *Macrosiphum euphorbiae* aphid samples feeding on four varieties of rose (red country rose variety, white bunchy rose variety, pink country rose variety and red tea rose variety) from four different locations in the Pune city and preparation of their morphological slides. Morphometric studies were conducted to trace any local morphological adaptations or existence of any host plant based populations supported by morphological adaptations.

Materials and Methods

A) Aphid collections

Individual aphids of *M. euphorbiae* were collected from different varieties of rose plants (*Rosa* sp.). They occur as green and pinkish brown in life colour. The host plant species include red country rose var., white bunchy rose var., pink country rose var. and red tea rose var. (flower having fragrance like tea). Adult apterae aphids were collected with insect brush and stored in 10 ml vials containing 70% alcohol. Aphid collection records of the adult apterous viviparous

Corresponding Author:

Parna Bhadra

Department of Zoology, MES

Abasaheb Garware College

(Autonomous), Pune,

Maharashtra, India

M. euphorbiae were kept while collections were cleared and mounted in Canada balsam on microscope slides.

B) Character choice

Ten standard morphological characters for aphids were scored [12, 13, 14]. Morphological measurements were made using a binocular transmission microscope fitted with a calibrated micrometer eyepiece [15]. Before the measurements were

made, the microscope magnification was calibrated and kept undisturbed during the study. Body length (BL), body width (BW), caudal length (CAU), siphunculi (SIPH), last antennal segment (PT or Processus Terminalis), base of antenna (B-ANT), rostrum (ROST), ultimate rostral segment (URS), head (HEAD), total antennal length (ANT) were the characters studied. For each locality, 20 apterous adult aphids were taken for measurements.



Macrosiphum euphorbiae in life colour.

The representation for the four populations were done by numbering as follows - the SB Road population (from red country var.) numbered as 1, the AGC population (from white

bunchy var.) numbered as 2, Camp population (from red tea var.) numbered as 3 and the Khadakwasla population (from country pink var.) was numbered as 4.




Fig 1: Characters of aphids dealt in this study (replicated from AFJ Dixon, 1994) [16]

C) Data analysis: The raw data was analysed through simple mean values and represented graphically

Results

The difference in morphometric data indicated the existence of four different locally adapted host plant based populations. Each of the ten characters was specifically distinguishable for

the host plant. The BL, SIPH, PT, URS, B-ANT and ROST was highest in the Khadakwasla population, BW was highest in both AGC and Khadakwasla population, ANT was seen to be longest in the Camp population. The SB Road population showed smallest size in morphometry in almost all the ten characters. All measurements were taken in millimeter scale (mm).






Fig 2: Comparison of the URS and ROST length for the four populations

**Fig 3:** Comparison of ANT, B-ANT and PT**Fig 4:** Comparison of SIPH and CAU length from the four populations

Discussion

Each of the four populations of *Macrosiphum euphorbiae* in our study showed some or the other phenotypic difference in characters. Morphometric analyses indicated the existence of four morphologically differentiated groups within *M. euphorbiae* associated with the rose plant varieties providing evidence for the presence of four host-associated groups. Insects studied on cottonwood plants, showed that similar genetic traits support similar communities [17]. The most important Characteristics observed in our study were the length of processus terminalis, siphunculus, rostrum and ultimate rostral segment. The siphunculus length in *M. euphorbiae* aphids in general is considerably larger than that of other aphid populations, a characteristic in which the population could be adapted to environmental conditions. The siphunculi are known to secrete alarm pheromones and probably aphids with longer siphunculi are able to release the alarm pheromone in a wider space [18]. Since the microclimatic conditions of that host plant are specific, all aphids and also *M. euphorbiae* are different from different host plants [19]. Rostral length on the other hand is associated with the ability of the aphid to reach deeper levels of phloem sap. Stronger rostrum enables them to penetrate tougher plant tissue. The key character is the ultimate rostral segment that has a layer of chitin enabling the rostrum to make the incision in the host plant tissue. It also resists the flow of phloem sap. The Khadakwasla and Camp populations had longer rostrum and ultimate rostral segment. The aphids sense their host plant tissue having mucilaginous cells and other chemical secretions with the antennae especially the processus terminalis. Longer antennae in the Camp population is an indication of the better adaptability of the population to sense the host tissue for the depth to which phloem is located. Further studies are to be done regarding the genetic variation of the host plants supporting the studied phloem feeding populations of the same aphid. The biological studies and ecological studies of the aphid are to be visited to support our current findings.

Acknowledgements

Appreciative thanks are due to the gardening staff of the MES Abasaheb Garware College, DIAT Park, Khadakwasla, Empress Botanical Garden, Camp and Parimal Udyan, S. B. Road for helping us during the study. We thank the non-teaching staff of the Department of Zoology, MES Abasaheb Garware College for having patience and helping us while carrying out morphological studies or waiting with us in the department late in the evening hours. Thanks are due to the teaching staff, especially Dr. G. A. Haval, HOD, Deptt. of Zoology, MES Abasaheb Garware College for encouragement and resource support in the department required for our study.

References

- Singh R, Singh G. Aphids. In: Omkar, editor. Polyphagous Pests of Crops. Singapore: Springer Nature; 2021. p.105-182.
- Blackman RL, Eastop VF. Aphids on the World's Crops: An Identification and Information Guide. London: Wiley & Sons; 2000.
- Dixon AFG. The Biology of Aphids. London: Edward Arnold; 1973.
- Blackman R, Eastop VF. Aphids on the World's Herbaceous Plants and Shrubs. Vol. 1: Host Lists and Keys; Vol. 2: The Aphids. London: John Wiley & Sons Ltd; 2007. 1439 p.
- Raychoudhuri DN, Ghosh D, Agarwala BK. Studies on the aphids from South India I. Insecta Matsumurana. 1981;23:1-20.
- Johnson MTJ. Bottom-up effects of plant genotype on aphids, ants, and predators. Ecology. 2008;89(1):145-154.
- Smith CM, Liu X, Wang LJ, Chen MS, Starkey S, Bai J. Aphid feeding activates expression of a transcriptome of oxylipin-based defense signals in wheat involved in resistance to herbivory. J Chem Ecol. 2010;36:260-276.
- Via S, Hawthorne JD. The genetic architecture of ecological specialization: correlated gene effects on host use and habitat choice in pea aphids. Am Nat. 2002;159(Suppl 3):S76-S88.
- Vorburger C. Temporal dynamics of genotypic diversity reveal strong clonal selection in the aphid *Myzus persicae*. J Evol Biol. 2006;19(1):97-107.
- Figueroa CC, Simon JC, Le Gallic JF, Leterme NP, Briones LM, Dedryver CA, Niemeyer HM. Effect of host defense chemicals on clonal distribution and performance of different genotypes of the cereal aphid Sitobion avenae. J Chem Ecol. 2004;30(12):2515-2525.
- Woolley VC, Tembo LB, Ndakidemi B, Obanyi JN, Arnold EJ, Belmain SR, Stevenson PC. The diversity of aphid parasitoids in East Africa and implications for biological control. Pest Manag Sci. 2022;78(3):1109-1116.
- Blackman RL, De Boise E. Morphometric correlates of karyotype and host plant in the genus *Euceraphis* (Hemiptera: Aphididae). Syst Entomol. 2002;27:323-335.
- Margaritopoulos JT, Tsourapas C, Tzortzi M, Kanavaki OM, Tsitsipis JA. Host selection by winged colonisers within the *Myzus persicae* group: a contribution towards understanding host specialization. Ecol Entomol. 2005;30:406-418.
- Bhadra P, Agarwala BK. A comparison of fitness characters of two host plant-based congeneric species of the banana aphid, *Pentalonia nigronervosa* and *P. caladii*. J Insect Sci. 2010;10(140):1-13.
- Raychoudhuri DN, Ghosh LK, Das SK. Studies on the aphids (Homoptera: Aphididae) from north and north-west India. Insecta Matsumurana. 1980;20:1-42.
- Dixon AFG, Kindlmann P. Optimum body size in aphids. Ecol Entomol. 1994;19(2):121-126.
- Barbour AG, Bunikis J, Travinsky B, Hoen AG, Diuk-Wasser MA, Fish D, Tsao JA. Niche partitioning of *Borrelia burgdorferi* and *Borrelia miyamotoi* in the same tick vector and mammalian reservoir species. Am J Trop Med Hyg. 2009;81(6):1120-1131.
- Dixon AFG. Insect Herbivore-Host Dynamics: Tree Dwelling Aphids. Cambridge: Cambridge University Press; 2004.
- Evans EW. Multitrophic interactions among plants, aphids, alternate prey and shared natural enemies: a review. Eur J Entomol. 2008;105(3):369-380.